

WHO/GPA/TCO/PMT/95.18

Distr.: General

Orig.: English

FLOW-CHARTS FOR SYNDROMIC CASE MANAGEMENT OF STD




WORLD
HEALTH
ORGANIZATION

© World Health Organization 1995

This document is not a formal publication of the World Health Organization (WHO), and all rights are reserved by the Organization. The document may, however, be freely reviewed, abstracted, reproduced or translated, in part or in whole, but not for sale or for use in conjunction with commercial purposes.


The views expressed in documents by named authors are solely the responsibility of those authors.

VAGINAL DISCHARGE




* **Positive** = age <21 years; or single; or > 1 partner; or new partner in past 3 months


URETHRAL DISCHARGE



GENITAL ULCERS



SCROTAL SWELLING



Patients complains of lower abdominal pain

LOWER ABDOMINAL PAIN

Take history and examine

Missed/overdue period or
Recent delivery/abortion or
Rebound tenderness or
Guarding or
Vaginal bleeding

NO

Temperature 38°C or
Pain during examination
or Vaginal discharge

NO

*Follow up if
pain persists*

YES

Refer

YES

- *Treat for PID*
- *Educate*
- *Counsel if needed*
- *Promote/provide condoms*
- *Partner management*

*Follow up after 3 days or
sooner if pain persists*

Improved?


NO

Refer

YES

Continue treatment

NEONATAL CONJUNCTIVITIS



INGUINAL BUBO

