

WHO/Mal/191 Add.1 ✓
WHO/Insecticides/68 Add.1
29 November 1957

ENGLISH ONLY

SUSCEPTIBILITY OF ADULT ANOPHELINES TO DDT AND OF CULEX FATIGANS
AND AÈDES AEGYPTI LARVAE TO DDT AND DIELDRIN

The following Editors' note which it was intended to add to the above paper is brought to the readers' attention:

It should be noted that in the present investigation the author has not entirely followed Busvine and Nash method because he expresses the concentration of DDT as the concentration in the oil-ether mixture (see table at page 2 and table 1, page 4). According to Busvine and Nash, inasmuch as the volatile solvent (trichloroethylene or ether) evaporates, the concentration to be stated is the final one, i.e., that of the insecticide in the Risella oil and not in the oil-ether mixture. Hence the results given in this paper should be re-elaborated¹ in order to make them comparable with those of other workers who follow strictly Busvine and Nash's method.

Further in this paper, in table 2 at page 5, the dosage is expressed in g/m^2 as if all the insecticide was on one surface of the filter paper, which obviously is not the case. Ed.

¹ Thus the table on page 2 should be replaced by the following:

Series	Mixture used		% DDT in 1 ml of mixed solution	% DDT in 0.33 ml of Risella oil solution
	Solution of DDT in Risella	Commercial ether		
B	2 ml at 2%	4 ml	0.66	2.0
C	2 ml at 1%	4 ml	0.33	1.0
Controls	2 ml -	4 ml	-	-

The LC_{50} values cited on page 3 should thus be amended, as follows:

	<u>Cited value</u>	<u>Correct value</u>
<u>A. albimanus</u>	0.24% DDT	0.7% DDT
<u>A. aquasalis</u>	0.33% DDT	1.0% DDT
<u>A. albitarsis</u>	0.39% DDT	1.1% DDT

Ed.