

a 63753



WHO/Mal/451 Corr.1
WHO/Vector Control/70)Corr.1
17 July 1964

ENGLISH ONLY

A LARGE-SCALE FIELD TRIAL WITH DICHLORVOS AS A RESIDUAL FUMIGANT
INSECTICIDE IN KANKIYA DISTRICT, NORTHERN NIGERIA

(Preliminary report)

by

Dr C. V. Foll and Dr C. P. Pant

Corrigenda

Page 2, paragraph 1, line 8: for "Sahoto" read "Sokoto".

Page 3, paragraph 2, line 4: for "A surrounding field of millet or guinea corn"
read "a surrounding tall fence (Zana) of millet or guinea corn stalks".

Page 3, paragraph 2, line 14: delete the words "guinea corn stalks".

Page 3, paragraph 2, line 15: for "and comprise" read "and they comprise".

Page 3, paragraph 3, line 4: for "maximal value" read "maximal morning value".

Page 4, paragraph 3, line 3: for "70 km²" read "466 km²".

Page 7, paragraph 3, line 3: for "P. malariae: 12.6%, P. ovale: 0.1%, and
P. falciparum + P. malariae: 1.5%" read "P. malariae: 1.5%, P. ovale: 0.1%,
and P. falciparum + P. malariae: 12.6%".

Page 10, last 14 lines: The two paragraphs from "In an attempt to see . . ." to
". . . and 'possibly infected' at the same period" should be deleted as they
refer to figures not reproduced in the preliminary report.

Page 13, paragraph 1, line 3: for "action" read "concentration".

Page 15, paragraph (iii), line 2: for "Table 29" read "Table 19".

Page 19: Acknowledgements, line 5: for "Wamki" read "Wambai".

Figure 7 - Infection rates, comparison area (sporozoites only): The vertical
scale - sporozoite rate should read "0.0, 1.0, 2.0, 3.0, 4.0, 5.0".