

# Complementary feeding

Report of the global consultation

Summary of guiding principles



World Health  
Organization





# Complementary feeding

## Report of the global consultation

convened jointly by  
the Department of Child and Adolescent Health and Development and  
the Department of Nutrition for Health and Development  
and

## Summary of guiding principles

for complementary feeding  
of the breastfed child



**World Health Organization**

WHO Library Cataloguing-in-Publication Data

*Complementary feeding: report of the global consultation, and summary of guiding principles for complementary feeding of the breastfed child.*

1. Infant nutrition 2. Guidelines 3. Breast feeding 4. Feeding behavior I. Global Consultation on Complementary Feeding (2001 : Geneva, Switzerland) II. Title: Summary of guiding principles for complementary feeding of the breastfed child.

ISBN 92 4 154614 X

(NLM Classification: QU 145.5)

**© World Health Organization 2002**

All rights reserved. Publications of the World Health Organization can be obtained from Marketing and Dissemination, World Health Organization, 20 Avenue Appia, 1211 Geneva 27, Switzerland (tel: +41 22 791 2476; fax: +41 22 791 4857; email: [bookorders@who.int](mailto:bookorders@who.int)). Requests for permission to reproduce or translate WHO publications – whether for sale or for noncommercial distribution – should be addressed to Publications, at the above address (fax: +41 22 791 4806; email: [permissions@who.int](mailto:permissions@who.int)).

The designations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted lines on maps represent approximate border lines for which there may not yet be full agreement.

The mention of specific companies or of certain manufacturers' products does not imply that they are endorsed or recommended by the World Health Organization in preference to others of a similar nature that are not mentioned. Errors and omissions excepted, the names of proprietary products are distinguished by initial capital letters.

The World Health Organization does not warrant that the information contained in this publication is complete and correct and shall not be liable for any damages incurred as a result of its use.

Typeset and printed in Geneva, Switzerland

# TABLE OF CONTENTS

FOREWORD	iii
REPORT OF THE GLOBAL CONSULTATION	1
Introduction	1
Discussion themes	2
Conclusions	3
Meeting energy and nutrient requirements	3
Ensuring availability of adequate complementary foods	4
Improving feeding behaviours	5
Strengthening programmatic approaches to improving complementary feeding	6
Recommendations	8
Closing statement	10
SUMMARY OF GUIDING PRINCIPLES	11
ANNEXES	13
Annex 1: List of participants	13
Annex 2: Meeting agenda	19
Annex 3: References	22
Annex 4: Lessons learned from large-scale behaviour change programmes to improve infant and young child feeding	23
Annex 5: Checklist of actions for successful complementary feeding	24



## FOREWORD

Nutrition's importance as a foundation for healthy development is often underestimated. Poor nutrition leads to ill-health and ill-health contributes to further deterioration in nutritional status. These effects are most dramatically observed in infants and young children, who bear the brunt of the onset of malnutrition and suffer the highest risk of disability and death associated with it. In 2001, 50–70% of the burden of diarrhoeal diseases, measles, malaria and lower respiratory infections was attributable to malnutrition.

The children who die represent only a small part of the total health burden due to nutritional deficiencies. Maternal malnutrition and inappropriate breastfeeding and complementary feeding represent major risks to the health and development of those children who survive. Deficiencies in the diet of vitamin A, iodine, iron and zinc are still widespread and are a common cause of excess morbidity and mortality. Over 50 million children under age five are wasted, and in low-income countries one in every three children suffers from stunted growth. Indeed, many children never reach this age. The effects of poor nutrition and stunting continue throughout life, contributing to poor school performance, reduced productivity, and impaired intellectual and social development.

Inappropriate feeding practices are a major cause of the onset of malnutrition in young children. Children who are not breastfed appropriately have repeated infections, grow less well, and are almost six times more likely to die by the age of one month than children who receive at least some breast milk. From six months onwards, when breast milk alone is no longer sufficient to meet all nutritional requirements, infants enter a particularly vulnerable period of complementary feeding during which they make a gradual transition to eating family foods. The incidence of malnutrition rises sharply during the period from 6 to 18 months of age in most countries, and the deficits acquired at this age are difficult to compensate for later in childhood.

During the past decade, there has been considerable progress in the implementation of interventions to improve breastfeeding practices. Clear recommendations and guidelines, combined with political commitment and increased allocation of resources, enabled many governments to establish programmes that combined the necessary actions to protect, promote and support breastfeeding. Consequently, improvements in breastfeeding rates have been demonstrated in various settings.

However, similar progress has not been made in the area of complementary feeding. While research and development have contributed to an expanding evidence base for recommendations on appropriate feeding and effective interventions for children after six months of age, translation of new knowledge into action has lagged behind.

To consider this gap and what could be done to fill it, WHO convened a global consultation on complementary feeding (Geneva, 11–13 December 2001). A group of scientists and programme managers was invited to review and update recommendations for appropriate complementary feeding, and to identify actions needed to accelerate programmatic efforts, including priorities for research and development of tools to plan and

implement interventions. Discussing issues relating to foods and feeding, they also considered the intricate links between maternal nutrition and appropriate breastfeeding and complementary feeding practices.

This report presents the proceedings of the consultation; it is meant to guide policy-makers and programme planners at all levels in taking appropriate action to give effect to the Global Strategy for Infant and Young Child Feeding,<sup>1</sup> which the World Health Assembly adopted in May 2002. It is hoped that the information will motivate all concerned parties to make the investments required to ensure that the nutritional needs of infants and young children are met worldwide. At the same time, it is hoped that the results will stimulate research and development to broaden the range of interventions to improve infant and young child feeding.

---

<sup>1</sup> WHO. *Global Strategy for Infant and Young Child Feeding*. Geneva: World Health Organization. WHA55/2002/REC/1 Annex 2, 2002

# REPORT OF THE GLOBAL CONSULTATION

## INTRODUCTION

Malnutrition is responsible, directly or indirectly, for over half of all childhood deaths. Infants and young children are at increased risk of malnutrition from six months of age onwards, when breast milk alone is no longer sufficient to meet all nutritional requirements and complementary feeding needs to be started. Complementary foods are often of lesser nutritional quality than breast milk. In addition, they are often given in insufficient amounts and, if given too early or too frequently, they displace breast milk. Gastric capacity limits the amount of food that a young child can consume during each meal. Repeated infections reduce appetite and increase the risk of inadequate intakes. Infants and young children need a caring adult or other responsible person who not only selects and offers appropriate foods but assists and encourages them to consume these foods in sufficient quantity.

Global recommendations for appropriate feeding of infants and young children are:

- Breastfeeding should start early, within one hour after birth.
- Breastfeeding should be exclusive for six months.
- Appropriate complementary feeding should start from the age of six months with continued breastfeeding up to two years or beyond.

Appropriate complementary feeding is:

- *timely* – meaning that foods are introduced when the need for energy and nutrients exceeds what can be provided through exclusive and frequent breastfeeding;
- *adequate* – meaning that foods provide sufficient energy, protein, and micronutrients to meet a growing child’s nutritional needs;
- *safe* – meaning that foods are hygienically stored and prepared, and fed with clean hands using clean utensils and not bottles and teats;
- *properly fed* – meaning that foods are given consistent with a child’s signals of appetite and satiety, and that meal frequency and feeding method – actively encouraging the child to consume sufficient food using fingers, spoon or self-feeding – are suitable for age.

Precise definitions of appropriate practices have been developed for breastfeeding. Certain determinants that affect the timely initiation, establishment and continuation of breastfeeding have also been identified in families, communities and the health system. These definitions and determinants have formed the basis of international and national guidelines on the protection, promotion and support of breastfeeding, and have resulted in intensive programmatic action to improve breastfeeding practices in many countries. In contrast, appropriate complementary feeding has been more difficult to define in precise operational terms. This has served as a barrier to taking effective action and tracking progress, for example in promoting and measuring improvements in specific behaviours.

The scientific review on complementary feeding<sup>1</sup> that WHO and UNICEF commissioned in 1998 provided age-specific guidance on nutritional requirements from complementary foods in healthy breastfed children. The review's conclusions have been translated into practical guidance for improving caregiver practices, notably in the feeding component of the WHO/UNICEF strategy for Integrated Management of Childhood Illness (IMCI),<sup>2</sup> and the WHO manual *Complementary feeding: family foods for breastfed children*.<sup>3</sup> Since the publication of the scientific review, new estimates have been made for energy and nutrient requirements for infants and young children<sup>4</sup>, warranting an update of previous recommendations.

Reducing childhood malnutrition requires a multi-sectoral approach that includes a variety of interventions to address its major causes. There is increasing evidence for the positive impact of feeding counselling on energy and nutrient intakes and growth in children less than two years of age<sup>5</sup>. To support changes in individual behaviour, supplemental interventions will be needed in many settings to ensure the availability and utilization of adequate micronutrient-rich complementary foods. Given the close link between maternal health and child health outcomes, in particular the contribution of low birth weight to childhood malnutrition, interventions should also address the health and nutrition of mothers. Against this background, WHO convened the consultation to:

- update global recommendations for complementary feeding; and
- provide more precise guidance for programme implementers to take these recommendations into action.

## DISCUSSION THEMES

The consultation brought together over 60 experts from a variety of disciplines and agencies (Annex 1). Using information in five background papers<sup>6</sup> and various presentations (Annexes 2 and 3), participants reviewed up-to-date information on:

---

<sup>1</sup> WHO/UNICEF. *Complementary feeding of young children in developing countries: A review of current knowledge*. WHO/NUT/98.1 Geneva: World Health Organization, 1998.

<sup>2</sup> WHO/UNICEF. *Integrated Management of Childhood Illness: a training course for first-level health workers*. WHO/CHD/97.3.A–L. Geneva: World Health Organization, 1997

<sup>3</sup> WHO. *Complementary feeding: family foods for breastfed children*. WHO/NHD/00.1, WHO/FCH/CAH/00.6 Geneva: World Health Organization, 2000

<sup>4</sup> Dewey KG, Brown KH. Update on technical issues concerning complementary feeding of young children in developing countries and implications for intervention programs. *Food and Nutrition Bulletin*, 2003;24(1): 5–28

<sup>5</sup> Santos I, Victora CG, Martines J et al. Nutrition counselling increases weight gain among Brazilian children. *Journal of Nutrition*, 2001, 131 (11): 2866–2873.

<sup>6</sup> The background papers are published in the *Special Issue of the Food and Nutrition Bulletin*, 2003;24 (1). For references, see Annex 3.

- energy and nutrient requirements of infants and young children, and the relative contribution of complementary foods to meeting these needs in healthy breastfed children;
- appropriate caregiver behaviours for complementary feeding and factors influencing them;
- approaches for improving the quality of complementary foods at household and community levels;
- macro-level approaches for ensuring the availability and utilization of adequate complementary foods;
- lessons learned from successful programmes to improve complementary feeding.

Based on the information presented in the background papers and presentations, and on the results of group work, participants formulated conclusions and recommendations regarding appropriate feeding practices and programmatic approaches for their promotion.

## CONCLUSIONS

### *Meeting energy and nutrient requirements*

1. New data on energy requirements in infancy endorsed by a FAO/WHO/UNU consultation warrant a revision of recommendations presented in the WHO/UNICEF scientific review on complementary feeding published in 1998.

The latest estimated energy requirements from complementary foods, assuming an average breast-milk intake, are 200 kcal/day for infants aged 6–8 months, 300 kcal/day for infants aged 9–11 months, and 550 kcal/day for children aged 12–23 months.

2. Scientific and empirical data are not yet sufficiently robust to warrant a change in estimated nutrient requirements from complementary foods described in the scientific review. However, a comparison of average nutrient intakes of children aged 6–24 months and new dietary reference intakes published by the Institute of Medicine, USA shows that the diets of infants and young children in most populations in low-income countries are consistently deficient in some nutrients, including iron, zinc and vitamin B6.
3. Adequate energy and nutrient intakes for this age-group are the result of a balance between appropriate breastfeeding and complementary feeding. There is no evidence of a preferential order between breastfeeding and complementary feeding at a given meal. Support for sustained breastfeeding as part of efforts to improve complementary feeding is critical. Increasing complementary feeding frequency, for example, may impair breastmilk intake with the potential risk of reducing total energy and nutrient intake if not enough attention is paid to sustaining breastfeeding.

4. On a population basis, recommended meal frequencies – assuming a diet with energy density of 0.8 kcal per gram or above and low breastmilk intake – are:
  - 2–3 meals per day for infants aged 6–8 months;
  - 3–4 meals per day for infants aged 9–11 months and children 12–24 months
  - additional nutritious snacks may be offered 1–2 times a day, as desired.
5. Complementary foods should be varied and include adequate quantities of meat, poultry, fish or eggs, as well as vitamin A-rich fruits and vegetables every day. Where this is not possible, the use of fortified complementary foods and vitamin mineral supplements may be necessary to ensure adequacy of particular nutrient intakes.

As infants grow, the consistency of complementary foods should change from semi-solid to solid foods and the variety of foods offered should increase. By eight months, infants can eat ‘finger foods’ and by 12 months, most children can eat the same types of food as the rest of the family.

### *Ensuring availability of adequate complementary foods*

6. Access to adequate complementary foods is a necessary condition for improving infant and young child feeding. There is ample evidence that diets, if based solely on unmodified, locally available ingredients, are often inadequate to meet recommended energy and nutrient needs.
7. Approaches to improving the availability of adequate complementary foods include simple technologies that can be applied in the home or community, and larger-scale industrial production of fortified processed foods that can involve both the public and the private sector.
8. Household technologies such as fermentation, soaking, roasting and malting are traditionally used in many societies. They can contribute to improving the safety and quality of complementary foods. However, reaching an adequate nutrient level remains a concern, particularly in diets that are mainly plant-based.

Specific additional measures may be needed to enrich home-prepared complementary foods. Mixing of micronutrient sprinkles or nutrient-rich pastes has been successfully promoted in some settings, and requires further research. Regular supplementation with micronutrient drops or capsules is another option to fill certain dietary gaps, but it is programmatically challenging to reach coverage.

9. Rapid urbanization and changing social networks affect caregivers’ ability to use freshly prepared home-grown foods. Centrally processed fortified foods, which can play an important role in ensuring adequate complementary diets, have been successfully promoted in various settings.

Public-private partnerships can play an important role in making available nutritionally adequate low-cost processed foods. Such partnerships should be consistent with the

provisions in the International Code of Marketing of Breastmilk and subsequent relevant World Health Assembly resolutions, and applicable Codex Alimentarius standards.

Voluntary third-party certification is a way to stimulate healthy competition among producers, while ensuring quality of products. It requires the establishment of an independent quality control body that certifies products as being suitable for their purpose, as well as promotion of the quality seal to the general public.

### *Improving feeding behaviours*

10. Improving complementary feeding requires attention to foods as well as to feeding behaviour of caregivers. Infants and young children need assistance that is appropriate for their age and developmental needs to ensure that they consume adequate amounts of complementary food. This is called responsive feeding.

Critical dimensions of responsive feeding are:

- feeding with a balance between giving assistance and encouraging self-feeding, as appropriate to the child's level of development;
- feeding with positive verbal encouragement, without verbal or physical coercion;
- feeding with age-appropriate and culturally appropriate eating utensils;
- feeding in response to early hunger cues;
- feeding in a protected and comfortable environment.
- feeding by an individual with whom the child has a positive emotional relationship and who is aware of and sensitive to the individual child's characteristics, including changes in physical and emotional state.

Inappropriate feeding behaviours are an important determinant of malnutrition. Caregivers often are unaware of the importance of responsive feeding, or do not know how to practise it. They need support from health professionals and community-based workers to acquire the necessary knowledge and skills.

11. Feeding behaviours are anchored in a wider belief system that influences what, when, where and how people feed their children. The most effective interventions are based on an in-depth assessment of this system; they address major barriers, using various channels and resources to support behaviour change.

Current knowledge emphasizes the importance of focusing on the family rather than on individual caregivers in designing interventions to improve complementary feeding. Assessing time allocation and time constraints in relation to food preparation and feeding are critical, as is estimating the real costs associated with implementing new feeding recommendations.

12. It is also important to promote safe preparation, feeding and storage of complementary foods in efforts to improve complementary feeding. Contamination and the proliferation of pathogens in food are major underlying causes of childhood diarrhoea.

*Strengthening programmatic approaches to improving complementary feeding*

13. In spite of the critical importance of appropriate complementary feeding for child growth and development, interventions to improve feeding practices in children 6–24 months of age have received insufficient attention. The Global Strategy for Infant and Young Child Feeding provides a framework for action and calls for concerted efforts to improve complementary feeding together with support for sustained breastfeeding. It stimulates governments, involving all concerned parties, to:
  - adopt a comprehensive national policy on infant and young child feeding, in the context of national policies and programmes for nutrition, child and reproductive health, and related areas;
  - constitute a national body to lead the implementation of the strategy as a coordinated national response;
  - ensure that the health and other sectors are able to protect, promote and support appropriate infant and young child feeding practices;
  - adopt appropriate measures to implement the International Code of Marketing of Breastmilk Substitutes and relevant subsequent World Health Assembly resolutions;
  - enact imaginative legislation to protect the breastfeeding rights of working women.
14. Improving complementary feeding requires a multi-dimensional approach focusing on improved feeding practices, control of childhood diseases and maternal nutrition. When interventions include an emphasis on breastfeeding, particularly exclusive breastfeeding for the first six months, in addition to improved complementary foods, a growth effect is more likely to be observed. Thus, comprehensive approaches that address the full range of infant and young child feeding practices are needed. Disease prevention and control interventions need to be packaged with nutrition interventions to prevent that repeated infections reduce the growth benefits.
15. The effect of interventions to improve feeding practices is variable and depends on the initial nutritional status of the infants and the degree to which other health needs are addressed. Consequently, other measures beyond weight gain or linear growth should be included in measuring the impact of complementary feeding interventions, such as biochemical measures of micronutrient status, changes in morbidity, and neurobehavioral development.
16. There is ample opportunity in current child health programmes and initiatives to improve support for infant and young child feeding by integrating appropriate guidance into the delivery of care. For example, Integrated Management of Childhood Illness, Early Childhood Development, Baby-friendly Hospital Initiative, Expanded Programme for Immunization, and Prevention and Control of HIV/AIDS can all make substantive contributions to the implementation of the Global Strategy for Infant and Young Child Feeding.

An important example of an integrated strategy is the WHO/UNICEF Integrated Management of Childhood Illness which includes interventions that combine the management of common childhood illnesses with individual counselling of caregivers to adopt appropriate child feeding practices. IMCI is effective in improving infant and young child feeding. Its implementation also provides a model for coordination among programmes and sectors towards a common goal.

The essential nutrition actions approach is another useful example of integration. The definition of seven essential nutrition actions and six points of delivery<sup>1</sup> has guided programme managers in designing and implementing a minimum package of activities to improve nutrition in childhood.

17. Policies and programmes to improve complementary feeding require involvement and coordination across sectors, including health, nutrition, agricultural and social welfare. The private sector can also play an important role in improving access to and utilization of safe and adequate complementary foods.
18. Policies and programmes need to respond to the special needs of working women, by ensuring provision of effective maternity entitlements and facilities for working women to sustain breastfeeding.
19. Important lessons can be learned from programmes that were successful in improving infant and young child feeding. Useful points to keep in mind when planning interventions to improve infant and young child feeding are summarized in Annex 4 *Lessons learned from large-scale behaviour change programmes to improve infant and young child feeding*. Similarly, the *Checklist of actions for successful complementary feeding* in Annex 5 provides concrete guidance for staff in health facility and community-based programmes analogous to the *Ten steps to successful breastfeeding*.<sup>2</sup>
20. Investments in the promotion of improved infant and young child feeding need adequate monitoring and evaluation. While a set of global indicators to assess breastfeeding practices was agreed a decade ago and has since been used to assess progress, the absence of indicators for assessing complementary feeding practices has been a barrier to effective programme development.

---

<sup>1</sup> The essential nutrition actions (ENA) approach is based on taking key nutrition actions associated with improved health outcomes at the most relevant points of health service delivery contact. The seven ENA points include: 1 exclusive breastfeeding from birth to six months; 2. appropriate complementary feeding starting at six months along with breastfeeding up to 24 months or beyond; 3. appropriate feeding of infants and young children during and after illness; 4. adequate nutrition of women; 5. control of vitamin A deficiency; 6. control of anaemia through iron supplements and de-worming of women and children; and 7. control of iodine deficiency disorder. The six key points of health delivery contact include: 1. antenatal care; 2. delivery; 3. post-natal care/family planning; 4. immunization; 5. well-baby clinic; and 6. sick child visits.

<sup>2</sup> WHO. *Protecting, promoting and supporting breastfeeding: a joint WHO/UNICEF statement*. Geneva: World Health Organization, 1989

## RECOMMENDATIONS

1. The existing evidence supports moving forward rapidly with programmatic action to improve complementary feeding. The Global Strategy for Infant and Young Child Feeding provides the framework for governments and other concerned parties to fulfill their respective obligations and responsibilities. Given the fundamental importance of childhood nutrition as a basis for healthy development of individuals and nations, the implementation of the strategy should rank among the top public health priorities to be pursued in every country.
2. The *Guiding principles for complementary feeding of the breastfed child*,<sup>1</sup> endorsed by the participants, should be used by programme planners and implementers as a guide to develop locally appropriate feeding recommendations.
3. Complementary feeding is part of a continuum of linked intervention areas, which include maternal nutrition, breastfeeding, micro-nutrient supplementation, psycho-social stimulation, feeding and care during and after illness, and illness prevention and control. Leadership, guidance, coordination and implementation of infant and young child feeding programmes should come from groups best positioned to deal with this cluster of interventions.
4. In promoting improved complementary feeding, all concerned parties should adhere to the provisions of the International Code of Marketing of Breastmilk Substitutes and relevant subsequent World Health Assembly resolutions. The Codex Alimentarius is another important instrument for governments to ensure appropriate and consistent standards for processed complementary foods.
5. To broaden the range of effective interventions and programmatic approaches to improve complementary feeding, further research is needed in priority areas including:
  - estimating energy and nutrient requirements of children living in especially vulnerable circumstances, such as infants of HIV-positive mothers who are not breastfed, preterm and low birth weight infants;
  - assessing the effects of variations in energy density, feeding frequency, food quantity and food variety on total energy and nutrient intake, including intake of breast milk;
  - determining the optimal amount and type of lipid and fiber intake by children;
  - determining the bioavailability of iron and zinc in locally available foods;
  - assessing efficacy and effectiveness of fortified complementary foods, sprinkles and spreads in addressing dietary gaps, including optimal levels of formulations and ration sizes to improve nutrient intakes;
  - identifying factors affecting children's appetite and appropriate management of anorexia;

---

<sup>1</sup> Pan American Health Organization, *Guiding Principles for Complementary Feeding of the Breastfed Child*. Washington DC: Pan American Health Organization, World Health Organization, 2003. For a summary, see page 11.

- determining methods and criteria for characterizing the responsiveness of feeding styles in various settings and identifying effective methods for promoting responsive feeding;
  - determining the impact of improved responsive feeding on child growth and developmental outcomes;
  - developing strategies for sustaining breastfeeding as complementary foods are introduced and the young child progresses to the family diet;
  - developing and testing appropriate and effective strategies to ensure safe storage, preparation and feeding of complementary foods for infants and young children;
  - examining the use of linear programming for developing context-specific complementary feeding guidelines;
  - identifying alternative approaches for demand-creation of affordable and effective processed food products;
  - developing and testing indicators and tools for assessing complementary feeding practices, for use in programmes and global monitoring of trends.
6. There is an immediate need to harmonize estimates of nutrient requirements for infants and young children, and to develop precise guidelines for feeding infants and young children who are not breastfed.
7. Immediate actions that governments and international partners can take to move forward include:

*At the global level:*

- develop guidelines that describes the continuum of essential interventions to improve maternal nutrition, breastfeeding and complementary feeding practices, consistent with the Global Strategy for Infant and Young Child Feeding;
- agree on common definitions and indicators for appropriate complementary feeding;
- develop a clear rationale for investing in complementary feeding interventions based on existing evidence;
- develop detailed guidance for programme planners to plan and implement complementary feeding interventions, building upon key elements that have been identified as critical for inclusion in successful, large-scale programmes;
- move complementary feeding issues onto national and global agendas and incorporate actions into relevant programmes, including for agriculture, education, nutrition, health, and poverty alleviation.

*At the national level:*

- define a national policy on infant and young child feeding and care, with a detailed plan of action to accompany it;
- increase advocacy for complementary feeding and incorporate actions into relevant programmes, including agriculture, education, nutrition, health and poverty alleviation;
- develop a resource plan to obtain adequate human, financial and organizational resources;

- develop locally appropriate feeding recommendations and programmatic guidelines for their implementation, taking into account the needs of diverse communities and the range of available opportunities for improving infant and young child feeding and nutrition.

## **CLOSING STATEMENT**

The first two years of a child's life are a critical window during which the foundations for healthy growth and development are built. Infant and young child feeding is a core dimension of care in this period. Internationally agreed goals for reduction of malnutrition and child mortality will be achieved only if families receive the support they need to adequately care for their children, including their nutritional needs. This consultation has shown that evidence for effective interventions to support children's nutritional well-being is available and accumulating rapidly. The challenge now is to translate knowledge into action.

## SUMMARY OF GUIDING PRINCIPLES<sup>1</sup>

1. **DURATION OF EXCLUSIVE BREASTFEEDING AND AGE OF INTRODUCTION OF COMPLEMENTARY FOODS:** Practice exclusive breastfeeding from birth to six months of age, and introduce complementary foods at six months of age (180 days) while continuing to breastfeed.
2. **MAINTENANCE OF BREASTFEEDING:** Continue frequent, on-demand breastfeeding until two years of age or beyond.
3. **RESPONSIVE FEEDING:** Practice responsive feeding, applying the principles of psycho-social care. Specifically:
  - feed infants directly and assist older children when they feed themselves, being sensitive to their hunger and satiety cues;
  - feed slowly and patiently, and encourage children to eat, but do not force them;
  - if children refuse many foods, experiment with different food combinations, tastes, textures and methods of encouragement;
  - minimize distractions during meals if the child loses interest easily;
  - remember that feeding times are periods of learning and love – talk to children during feeding, with eye to eye contact.
4. **SAFE PREPARATION AND STORAGE OF COMPLEMENTARY FOODS:** Practice good hygiene and proper food handling by :
  - washing caregivers’ and children’s hands before food preparation and eating;
  - storing foods safely and serving foods immediately after preparation;
  - using clean utensils to prepare and serve food;
  - using clean cups and bowls when feeding children;
  - avoiding the use of feeding bottles, which are difficult to keep clean.
5. **AMOUNT OF COMPLEMENTARY FOOD NEEDED:** Start at six months of age with small amounts of food and increase the quantity as the child gets older, while maintaining frequent breastfeeding. The energy needs from complementary foods for infants with “average” breast milk intake in developing countries are approximately 200 kcal per day at 6–8 months of age, 300 kcal per day at 9–11 months of age, and 550 kcal per day at 12–23 months of age. In industrialized countries these estimates differ somewhat (130, 310 and 580 kcal/d at 6–8, 9–11 and 12–23 months, respectively) because of differences in average breast milk intake.

---

<sup>1</sup> Pan American Health Organization. *Guiding Principles for Complementary Feeding of the Breastfed Child*. Washington DC: Pan American Health Organization, World Health Organization, 2003

6. **FOOD CONSISTENCY:** Gradually increase food consistency and variety as the infant gets older, adapting to the infant's requirements and abilities. Infants can eat pureed, mashed and semi-solid foods beginning at six months. By eight months most infants can also eat "finger foods" (snacks that can be eaten by children alone). By 12 months, most children can eat the same types of foods as consumed by the rest of the family (keeping in mind the need for nutrient-dense foods, as explained in 8. below). Avoid foods that may cause choking (i.e., items that have a shape and/or consistency that may cause them to become lodged in the trachea, such as nuts, grapes, raw carrots).
7. **MEAL FREQUENCY AND ENERGY DENSITY:** Increase the number of times that the child is fed complementary foods as he/she gets older. The appropriate number of feedings depends on the energy density of the local foods and the usual amounts consumed at each feeding. For the average healthy breastfed infant, meals of complementary foods should be provided 2–3 times per day at 6–8 months of age and 3–4 times per day at 9–11 and 12–24 months of age. Additional nutritious snacks (such as a piece of fruit or bread or chapatti with nut paste) may be offered 1-2 times per day, as desired. Snacks are defined as foods eaten between meals, usually self-fed, convenient and easy to prepare. If energy density or amount of food per meal is low, or the child is no longer breastfed, more frequent meals may be required.
8. **NUTRIENT CONTENT OF COMPLEMENTARY FOODS:** Feed a variety of foods to ensure that nutrient needs are met. Meat, poultry, fish or eggs should be eaten daily, or as often as possible. Vegetarian diets cannot meet nutrient needs at this age unless nutrient supplements or fortified products are used (see 9. below). Vitamin A-rich fruits and vegetables should be eaten daily. Provide diets with adequate fat content. Avoid giving drinks with low nutrient value, such as tea, coffee and sugary drinks such as soda. Limit the amount of juice offered so as to avoid displacing more nutrient-rich foods.
9. **USE OF VITAMIN-MINERAL SUPPLEMENTS OR FORTIFIED PRODUCTS FOR INFANT AND MOTHER:** Use fortified complementary foods or vitamin-mineral supplements for the infant, as needed. In some populations, breastfeeding mothers may also need vitamin-mineral supplements or fortified products, both for their own health and to ensure normal concentrations of certain nutrients (particularly vitamins) in their breast milk. [Such products may also be beneficial for pre-pregnant and pregnant women.]
10. **FEEDING DURING AND AFTER ILLNESS:** Increase fluid intake during illness, including more frequent breastfeeding, and encourage the child to eat soft, varied, appetizing, favorite foods. After illness, give food more often than usual and encourage the child to eat more.

# ANNEXES

## Annex 1

### LIST OF PARTICIPANTS

GLOBAL CONSULTATION ON COMPLEMENTARY FEEDING  
10-13 DECEMBER 2001, GENEVA, ROOM D

**Cecilia Acuin**

Department of Clinical Epidemiology  
National Institute of Health  
University of the Philippines  
Manila, Philippines  
Tel: (632) 525 4098  
Fax: (632) 525 4098  
e-mail: cesacuin@pworld.net.ph

**Nita Bhandari**

All India Institute of Medical Science  
(AIMMS)  
Ansari Nagar  
New Delhi 110029, India  
Tel: (91) 11 601 4136  
Fax: (91) 11 601 6449  
e-mail: community.research@cih.uib.no

**Dure-Samin Akram**

Nutrition Support Programme  
Department of Pediatrics  
Unit 1, Civil Hospital  
Karachi, Pakistan  
Tel: (92) 21 921 5720  
Fax: (92) 21 921 5720  
e-mail: dsakram@gemini.khi.erum.com.pk

**Robert E. Black**

Department of International Health  
School of Hygiene and Public Health  
The Johns Hopkins University  
615 North Wolfe Street  
Baltimore, MD 21205-2179, USA  
Tel: (1) 404 955 3934  
Fax: (1) 404 955 1253  
e-mail: rblack@jhsp.edu

**Kunal Bagchi**

Regional Advisor, Nutrition  
WHO Regional Office for the Eastern  
Mediterranean  
Abdul Razzak Al Sanhoury Street  
Nasar City  
Cairo 11371, Egypt  
Tel: (202) 670 2534  
Fax: (202) 670 2492  
e-mail: bagchik@emro.who.int

**Ruth Bland**

Africa Centre for Population Studies and  
Reproductive Health  
P.O. Box 198  
Mtubatuba 3935  
KwaZulu Natal, South Africa  
Tel: (27) 35 550 0158  
Fax: (27) 35 550 1674  
e-mail: blandr@mrc.ac.za

**Geneviève Becker**

2 Kylemore Park  
Taylor's Hill  
Galway, Ireland  
Tel: (44) 353 91 527511  
Fax: (44) 353 91 528677  
e-mail: becker@iol.ie

**Neal Brandes**

Child Survival Division  
Office of Health and Nutrition  
USAID  
GH/HID/MCH 3.07/070  
Ronald Reagan Building  
1300 Pennsylvania Ave., N.W.  
Washington D.C. 20523, USA  
Tel: (1) 202 712 0771  
Fax: (1) 202 216 3702  
e-mail: nbrandes@usaid.gov

**Zuzana Brazdova**

Department of Preventive Medicine  
Masaryk University  
Jostova 10  
66244 Brno, Czech Republic  
Tel: (420) 602 578491  
Fax: (420) 542 126366  
e-mail: brazdova@med.muni.cz

**André Briend**

CNAM/IRD  
Institut Scientifique et Technique de la  
Nutrition et de l'Alimentation (ISTNA)  
5 rue du Vertbois  
75003 Paris, France  
Tel: (33) 1 53 01 80 36  
Fax: (33) 1 53 01 80 05  
e-mail: brienda@cnam.fr

**Kenneth Brown**

Programme in International Nutrition  
University of California  
One Shields Avenue  
Davis, CA 95616-8669, USA  
Tel: (1) 530 752 1992  
Fax: (1) 530 752 3406  
e-mail: khbrown@ucdavis.edu

**Reina Buijs**

Nutrition and Health  
Social Policy Division  
Ministry of Foreign Affairs  
Bezuidenhoutseweg 67  
2594 AC Den Haag, Netherlands  
Tel: (31) 70 348 5825  
Fax: (31) 70 348 5366  
e-mail: reina.buijs@minbuza.nl

**Laura Caulfield**

Center for Human Nutrition  
Department of International Health  
School of Hygiene and Public Health  
The Johns Hopkins University  
615 North Wolfe Street  
Baltimore, MD 21205-2179, USA  
Tel: (1) 410 955 2786  
Fax: (1) 410 955 0196  
e-mail: lcaulfie@jhsph.edu

**Tommaso Cavalli Sforza**

Regional Advisor, Nutrition  
WHO Regional Office for the Western  
Pacific  
P.O. Box 2932  
1099 Manila, Philippines  
Tel: (632) 52 88 001  
Fax: (632) 52 11 036  
e-mail: tommaso@wpro.who.int

**Visith Chavasit**

Institute of Nutrition  
Mahidol University  
Salaya, Phutthamonthon 4  
Nakhon Pathom 73170, Thailand  
Tel: (662) 800 2380 (ext. 416)  
Fax: (662) 441 9344  
e-mail: nuvca@mahidol.ac.th

**Hilary Creed de Kanashiro**

Instituto de Investigacion Nutricional  
Avenida La Molina  
685 La Molina  
Lima, Peru  
Tel: (51) 1 349 60 23  
Fax: (51) 1 349 60 25  
e-mail: hmcreed@iin.sld.pe

**Kathryn Dewey**

Department of Nutrition  
University of California  
Programme in International Nutrition  
University of California  
One Shields Avenue  
Davis, CA 95616-8669, USA  
Tel: (1) 530 752 1992  
Fax: (1) 530 752 3406  
e-mail: kgdewey@ucdavis.edu

**Tatang Sohibul Falah**

Complementary Feeding Programme  
Ministry of Health  
Jakarta, Indonesia  
Tel: (62) 021 527 7152  
Fax: (62) 021 521 0176  
e-mail: iodina@centrin.net.id

**Suzanne Farhoud**

Regional Advisor, Child Health  
 WHO Regional Office for the Eastern  
 Mediterranean  
 Abdul Razzak Al Sanhoury Street  
 Nasar City  
 Cairo 11371, Egypt  
 Tel: (202) 670 2534  
 Fax: (202) 670 2492  
 e-mail: farhouds@emro.who.int

**Rosalind Gibson**

Department of Human Nutrition  
 University of Otago  
 Dunedin, New Zealand  
 Tel: (64) 3 479 7955  
 Fax: (64) 3 479 7958  
 e-mail:  
 rosaling.gibson@stonebow.otago.ac.nz

**Boitshepo Giyose**

Commonwealth Institute  
 CRHCH-ECSA  
 P.O. Box 1009  
 Arusha, Tanzania  
 Tel: (255) 27 250 4105/6  
 Fax: (255) 27 250 4124/8292  
 e-mail: bgiyose@crhcs.or.tz

**Jaanaki Gooneratne**

Agro and Food Technology Division  
 Industrial Technology Institute  
 363, Baudhaloka Mawatha  
 Colombo 7, Sri Lanka  
 Tel: (94) 1 693 807  
 Fax: (94) 1 686 567  
 e-mail: jaanaki@iti.lk

**Agnes Guyon**

The Linkages Project  
 B.P. 5253 Antananarivo  
 Madagascar  
 Tel: (261) 20 22 613 52  
 Fax: (261) 20 22 613 52  
 e-mail: agnes.lkg@pact.mg

**Jean-Pierre Habicht**

Division of Nutritional Sciences  
 Cornell University  
 Ithaca, N.Y. 14853-6, USA  
 Tel: (1) 607 255 4419  
 Fax: (1) 607 255 2608  
 e-mail: jh48@cornell.edu

**Rukhsana Haider**

Nutrition Department  
 WHO Regional Office for South-East Asia  
 World Health House  
 Indraprastha Estate  
 Mahatma Gandhi Road  
 New Delhi 110002, India  
 e-mail: haiderr@whosea.org

**Iqbal Kabir**

International Centre for Diarrhoeal Disease  
 Research (ICDDR,B)  
 G.P.O. Box 128  
 Dhaka 1000, Bangladesh  
 Tel: (880) 2 881 1751-60  
 Fax: (880) 2 882 3116  
 e-mail: ikabir@icddr.org

**Wijnand Klaver**

Department of Agricultural Production and  
 Nutrition  
 International Agricultural Centre  
 P.O. Box 88  
 Wageningen, Netherlands  
 Tel: (31) 317 495366  
 Fax: (31) 317 495395  
 e-mail: w.klaver@iac.agro.nl

**Miriam Labbok**

Infant and Young Child Feeding  
 UNICEF  
 3 UN Plaza, Room 726  
 New York, N.Y. 10017, USA  
 Tel: (1) 212 824 6371  
 Fax: (1) 212 824 6465  
 e-mail: mlabbok@unicef.org

**Anna Lartey**

Department of Nutrition and Food Science  
 University of Ghana  
 P.O. Box LG134  
 Legon, Ghana, West Africa  
 Tel: (233) 21 513293  
 Fax: (233) 21 500389  
 e-mail: aalartey@hotmail.com

**Chessa Lutter**

Regional Advisor, Nutrition  
 WHO Regional Office for the Americas  
 525, 23<sup>rd</sup> Street, N.W.  
 Washington, D.C. 20037, USA  
 Tel: (1) 202 974 3000  
 Fax: (1) 202 974 3663  
 e-mail: lutterch@paho.org

**Elizabeth Mason**

Regional Advisor, Child Health  
 WHO Regional Office for Africa  
 Highlands Office  
 P.O. Box BE 773  
 Harare  
 Zimbabwe  
 Tel: (263) 4 746 127  
 Fax: (263) 4 746 127  
 e-mail: masone@whoafr.org

**Patience Mensah**

Noguchi Memorial Institute for Medical  
 Research  
 Legon, Ghana (on sabbatical at Oxford  
 University, UK)  
 c/o St. Hilda's College  
 Oxford, OX4 1DY, UK  
 Tel: (44) 1865 431040  
 Fax: (44) 1835 276816  
 e-mail: paciencemensah@st-  
 hildas.oxford.ac.uk

**Kim F. Michaelsen**

Research Department of Human Nutrition  
 Royal Veterinary and Agricultural University  
 Rolighedsvej 30  
 DK-1958 Copenhagen, Denmark  
 Tel: (45) 35 282493/95  
 Fax: (45) 35 282483  
 e-mail: kfm@kvl.dk

**Claire Mouquet**

Research Institute for Development  
 911 avenue Acropolis  
 34090 Montpellier, France  
 Tel: (33) 467 41 62 47  
 Fax: (33) 467 54 78 00  
 e-mail: mouquet@mpl.ird.fr

**Gretel Pelto**

Division of Nutritional Sciences  
 Cornell University  
 Ithaca, N.Y. 14853-6, USA  
 Tel: (1) 607 255 6277  
 Fax: (1) 607 255 2608  
 e-mail: gp32@cornell.edu

**Ellen Piwoz**

Centre for Nutrition  
 Academy for Education and Development  
 1825 Connecticut Avenue, N.W.  
 Washington D.C. 20009, USA  
 Tel: (1) 202 884 8816  
 Fax: (1) 202 884 8447  
 e-mail: epiwoz@aed.org

**Victoria Quinn**

The Linkages Project  
 1825 Connecticut Avenue, N.W.  
 Washington D.C. 20009-5721, USA  
 Tel: (1) 202 884 8829  
 Fax: (1) 202 884 8447  
 Email: vquinn@aed.org

**Sonya Rabeneck**

United Nations ACC/SCN Secretariat  
 Geneva, Switzerland  
 Tel: (41) 22 791 0456  
 Fax: (41) 22 798 8891  
 e-mail: rabenecks@who.int or  
 accsc@who.int

**Aileen Robertson**

Regional Advisor Nutrition  
 WHO Regional Office for Europe  
 8, Scherfigsvej  
 DK-2100 Copenhagen, Denmark  
 Tel: (45) 39 17 17 17  
 Fax: (45) 39 17 18 18  
 e-mail: aro@who.dk

**Claudia Rokx**

Health Nutrition and Population  
World Bank  
Room G7-043 (MSN G7-701)  
1818 H. Street, N.W.  
Washington D.C. 20433, USA  
Tel: (1) 202 458 2665  
Fax: (1) 202 522 3243  
e-mail: crokx@worldbank.org

**Marie Ruel**

Food Consumption and Nutrition Division  
IFPRI  
2033 K. Street, N.W.  
Washington D.C. 20006, USA  
Tel: (1) 202 862 5676  
Fax: (1) 202 467 4439  
e-mail: m.ruel@cgiar.org

**Tina Sanghvi**

USAID BASICS Project  
Suite 300, 1600 Wilson Boulevard  
Arlington, VA 22209, USA  
Tel: (1) 703 312 6893  
Fax: (1) 703 312 6900  
e-mail: tsanghiv@basics.org

**Ina Santos**

Department of Social Medicine  
University of Pelotas  
Av. Duque de Caxias, no 250  
Caixa postal 464  
Pelotas, Brazil  
Tel: (53) 271 2442  
Fax: (53) 271 2645  
e-mail: inasantos@uol.com.br

**Kirsten Simondon**

Research Institute for Development  
911 avenue Acropolis  
34090 Montpellier, France  
Tel: (33) 467 41 61 90  
Fax: (33) 467 54 78 00  
e-mail: kirsten.simondon@mpl.ird.fr

**Andrew Tomkins**

Centre for International Health  
Institute of Child Health  
30 Guilford Street  
London WC1 1EH, UK  
Tel: (44) 20 7905 2123  
Fax: (44) 20 7404 2062  
e-mail: a.tomkins@ich.ucl.ac.uk

**Kraisid Tontisirin**

Food and Nutrition Division  
Food and Agriculture Organization  
Viale della Terme di Caracalla  
00100 Rome, Italy  
Tel: (39) 6 5705 3330  
Fax: (39) 6 5705 4593  
e-mail: kraisid.tontisirin@fao.org

**Shakila Zaman**

Department of Social and Preventive  
Paediatrics  
King Edward Medical College  
Lahore, Pakistan  
Tel: (92) 12 723 3509  
Fax: (92) 12 723 3509  
e-mail: prevke12@lhr.paknet.com.pk

**Sheila Vir**

Nutrition Project  
UNICEF  
1/4 Vipul Khand  
Gomati Nagar  
Lucknow 226010, India  
Tel: (91) 522 303152-57  
Fax: (91) 522 303158  
e-mail: svir@unicef.org

**WHO Secretariat****Bruno de Benoist**

Department of Nutrition for Health and Development

Tel: (41) 22 791 3412

Fax: (41) 22 791 4156

e-mail: debenoistb@who.int

**Carmen Casanovas**

Department of Child and Adolescent Health and Development

Tel: (41) 22 791 4225

Fax: (41) 22 791 4853

e-mail: casanovasm@who.int

**Graeme Clugston**

Director, Department of Nutrition for Health and Development

Tel: (41) 22 791 3326

Fax: (41) 22 791 4156

e-mail: clugstong@who.int

**Bernadette Daelmans**

Department of Child and Adolescent Health and Development

Tel: (41) 22 791 2908

Fax: (41) 22 791 4853

e-mail: daelmansb@who.int

**Sultana Khanum**

Department of Nutrition for Health and Development

Tel: (41) 22 791 2624

Fax: (41) 22 791 4156

e-mail: khanums@who.int

**José Martines**

Department of Child and Adolescent Health and Development

Tel: (41) 22 791 2634

Fax: (41) 22 791 4853

e-mail: martinesj@who.int

**Mirella Mokbel-Genequand**

Department of Nutrition for Health and Development

Tel: (41) 22 791 2758

Fax: (41) 22 791 4156

e-mail: mokbelm@who.int

**Leda Nemer**

Department of Child and Adolescent Health and Development

Tel: (41) 22 791 3713

Fax: (41) 22 791 4853

e-mail: nemerl@who.int

**Randa Saadeh**

Department of Nutrition for Health and Development

Tel: (41) 22 791 3315

Fax: (41) 22 791 4156

e-mail: saadehr@who.int

**Jorgen Schlundt**

Department of Food Safety

Tel: (41) 22 791 3445

Fax: (41) 22 791 4807

e-mail: schlundtj@who.int

**Hans Troedsson**

Director, Department of Child and Adolescent Health and Development

Tel: (41) 22 791 3281

Fax: (41) 22 791 4853

e-mail: troedssonh@who.int

**Constanza Vallenias**

Department of Child and Adolescent Health and Development

Tel: (41) 22 791 4143

Fax: (41) 22 791 4853

e-mail: vallenasc@who.int

**Jelka Zupan**

Department of Reproductive Health and Research

Tel: (41) 22 791 4221

Fax: (41) 22 791 4171

e-mail: zupanj@who.int

## AGENDA

### GLOBAL CONSULTATION ON COMPLEMENTARY FEEDING

10-13 DECEMBER 2001, GENEVA, ROOM D

#### Monday, 10 December

9.00 – 9.30    Opening  
                  Introduction of participants  
                  Objectives of the meeting  
                  Expected outcomes

#### Technical update on issues related to access to adequate and safe complementary foods

9.30 - 10.30    Update on current scientific knowledge regarding energy, protein and micro-nutrients required from complementary foods (Dr Kenneth Brown, Dr Kathryn Dewey)

Discussion

10.30 - 11.00    Tea/coffee

11.00 - 12.00    Approaches for improving the availability of adequate complementary foods on large scale, including conclusions of the technical consultation on fortification of complementary foods (Dr Chessa Lutter)

Discussion

12.00 - 13.00    Household and community-based approaches to improve availability of safe and adequate complementary foods (Dr Rosalind Gibson, Dr Patience Mensah)

Discussion

13.00            Lunch

#### Technical update of issues related to improving feeding practices

14.00 - 15.00    Improving feeding practices: current patterns, common constraints, and the design of interventions in relation to best practices (Dr Gretel Pelto)

Discussion

15.00 - 15.45 Ensuring safe preparation and storage of foods (Dr Jurgen Schlundt)

Discussion

15.45 Tea/coffee

16.00 Introduction to group work

16.15 -17.30 Participants work in three groups

## **Tuesday, 11 December**

### Issues related to programmatic approaches to promote optimal complementary feeding

9.00 - 10.00 Components of successful complementary feeding programmes  
(Dr Kathryn Dewey)

Discussion

10.00 -10.30 Implementation of interventions to improve complementary feeding:  
case- study from India (Dr Nita Bhandari)

10.30 Tea/coffee

11.00 -12.00 Implementation of interventions to improve complementary feeding:  
case study from Peru (Dr Hilary Creed de Kanashiro)

Discussion of case studies

12.00 - 13.00 Experience with implementation of large-scale nutrition interventions  
(Ms Claudia Rokx)

Discussion

13.00 Lunch

14.00 - 15.00 Assessing complementary feeding practices: towards development of  
indicators (Dr Marie Ruel)

Discussion

15.00 - 17.30 Group work (continued)

**Wednesday, 12 December**

9.00 - 10.00 Challenges in promoting interventions to improve complementary feeding: lessons learned and possible solutions (Dr Ellen Piwoz)

Discussion

10.00 - 12.30 Groups develop recommendations and prepare their presentation

12.30 Lunch session: Implementation of interventions to improve infant and young child feeding: a case study from Madagascar (Dr Agnes Guyon)

13.30 - 15.30 Presentation of Group 1: Improving access to adequate and safe complementary foods

Discussion

15.30 - 16.00 Tea/coffee

16.00 - 18.00 Presentation of Group 2: Improving feeding practices

Discussion

**Thursday, 13 December**

9.00 - 10.30 Presentation of Group 3: Strengthening programmes to promote optimal complementary feeding

Discussion

10.30 Tea/coffee

11.00 - 12.30 Groups finalize their conclusions and recommendations and develop a summary of recommended priority activities

12.30 Lunch

14.00 - 16.45 Presentation of conclusions, recommendations and priority activities

- Improving access to adequate and safe complementary foods
- Improving feeding practices
- Strengthening programmes to promote optimal complementary feeding

16.45 Closing

## Annex 3

**REFERENCES**

## BACKGROUND PAPERS DEVELOPED IN PREPARATION OF THE CONSULTATION

Dewey KG, Brown KH. Update on technical issues concerning complementary feeding of young children in developing countries and implications for intervention programs. *Food and Nutrition Bulletin*, 2003;24(1): 5–28

Piwoz GE, Huffmann SL, Quinn VJ. Promotion and advocacy for improved complementary feeding: Can we apply the lessons learned from breastfeeding? *Food and Nutrition Bulletin*, 2003;24 (1): 29–44

Pelto GH, Levitt E, Thiaru L. Improving feeding practices: current patterns, common constraints, and the design of interventions. *Food and Nutrition Bulletin*, 2003;24(1): 45–82

Lutter CK. Macro-level approaches to improve the availability of complementary foods. *Food and Nutrition Bulletin*, 2003;24(1): 83–103

Mensah P, Tomkins A. Household-level technologies to improve the availability and preparation of adequate and safe complementary foods. *Food and Nutrition Bulletin*, 2003;24(1):104–125

The full version of the Special Issue of the Food and Nutrition Bulletin 24; (1) 2003 is available at

[http://www.who.int/child-adolescent-health/New\\_Publications/NUTRITION/FNB\\_24-1\\_WHO.pdf](http://www.who.int/child-adolescent-health/New_Publications/NUTRITION/FNB_24-1_WHO.pdf)

**Annex 4****LESSONS LEARNED FROM LARGE-SCALE BEHAVIOUR CHANGE PROGRAMMES TO IMPROVE INFANT AND YOUNG CHILD FEEDING**

- The development of a comprehensive policy and strategy for infant and young child feeding is critical to support programme actions. This policy should be based on international guidelines and adapted to local circumstances; sufficient resources should be invested in advocacy for its implementation.
- Programme action should be based on local adaptation of generic guidelines, using research to identify appropriate feeding recommendations together with interventions to facilitate their adoption.
- Based on feeding recommendations thus identified, a feasible package of desired behaviours should be designed for caregivers and families. Families should be provided skilled support to develop the skills they need in order to practice these behaviours effectively.
- In addition to caregivers, other family members should be targeted, since it is the family which makes decisions concerning feeding of infants and young children.
- The importance of negotiation should be emphasized in the interaction between a feeding counsellor and a caregiver or family. Interventions should work towards ‘incremental’ changes in feeding practices: changes should be small and practical, made step-by-step over time.
- Interventions should address a variety of actors that are part of the support network of families, so that consistent messages on infant and young child feeding are given at all levels, in the public and private health system, the media, and in communities.
- Critical knowledge and skills in support of infant and young child feeding should be incorporated into the pre-service training of health and other development professionals, so that sustainable implementation can be ensured.

**Annex 5****CHECKLIST OF ACTIONS FOR SUCCESSFUL COMPLEMENTARY FEEDING<sup>1</sup>**

Every facility and community programme involved in providing care for children 6–24 months of age should:

1. Have a written policy that is communicated to all health-care staff about complementary feeding.
2. Develop local feeding guidelines based on formative research.
3. Train all health-care and community workers in skills necessary to implement the policy and guidelines.
4. Inform all caregivers and parents about the benefits and management of complementary feeding.
5. Help mothers initiate complementary feeding at six months, while giving adequate support to sustain breastfeeding.
6. Show mothers how to safely prepare and offer complementary foods, while maintaining and supporting breastfeeding, according to the age and circumstances of the child.
7. Teach mothers about feeding frequency, food variety, and adequate quantities for growing children.
8. Counsel mothers how to maintain adequate feeding during and following illness and loss of appetite.
9. Establish infant feeding and care support groups and refer mothers to them.
10. Refer all mothers and children who are malnourished, sick, or living in families with special circumstances to health care and available family support services.

---

<sup>1</sup> Piwoz GE, Huffmann SL, Quinn VJ. Promotion and advocacy for improved complementary feeding: Can we apply the lessons learned from breastfeeding? *Food and Nutrition Bulletin*, 2003;24 (1):29–44.

## SELECTED WHO PUBLICATIONS OF RELATED INTEREST

WHO/UNICEF/Orstrom/University of California at Davis. *Complementary feeding of young children in developing countries*. WHO/NUT/98.1. Geneva: World Health Organization, 1998  
[http://www.who.int/child-adolescent-health/publications/NUTRITION/WHO\\_NUT\\_98.1.htm](http://www.who.int/child-adolescent-health/publications/NUTRITION/WHO_NUT_98.1.htm)

WHO. *The optimal duration of exclusive breastfeeding. A systematic review*. WHO/NHD/01.08 or WHO/FCH/CAH/01.23. Geneva: World Health Organization, 2002  
[http://www.who.int/child-adolescent-health/publications/NUTRITION/WHO\\_FCH\\_CAH\\_01.23.htm](http://www.who.int/child-adolescent-health/publications/NUTRITION/WHO_FCH_CAH_01.23.htm)

WHO. *Report of the Expert Consultation on the optimal duration of exclusive breastfeeding*. WHO/NHD/01.09 or WHO/FCH/CAH/01.24. Geneva: World Health Organization, 2002  
[http://www.who.int/child-adolescent-health/publications/NUTRITION/WHO\\_FCH\\_CAH\\_01.24.htm](http://www.who.int/child-adolescent-health/publications/NUTRITION/WHO_FCH_CAH_01.24.htm)

WHO. *Nutrient adequacy of exclusive breastfeeding for the term infant during the first six months of life*. Geneva: World Health Organization, 2002  
[http://www.who.int/child-adolescent-health/publications/NUTRITION/Nutrient\\_Adequacy.htm](http://www.who.int/child-adolescent-health/publications/NUTRITION/Nutrient_Adequacy.htm)

WHO. *Complementary feeding: Family foods for breastfed children*. WHO/NHD/00.1 or WHO/FCH/CAH/00.6. Geneva: World Health Organization, 2002  
[http://www.who.int/child-adolescent-health/publications/NUTRITION/WHO\\_FCH\\_CAH\\_00.6.htm](http://www.who.int/child-adolescent-health/publications/NUTRITION/WHO_FCH_CAH_00.6.htm)

WHO. *Global strategy for infant and young child feeding*. WHA55/2002/REC/1 Annex 2 or Geneva: World Health Organization, 2002.  
[http://www.who.int/gb/EB\\_WHA/PDF/WHA55/ea5515.pdf](http://www.who.int/gb/EB_WHA/PDF/WHA55/ea5515.pdf)

International Food Policy Research Institute/WHO. *Moving forward with complementary feeding: identifying indicators and research priorities*. Washington DC: FCND Discussion Paper 146, 2003  
[http://www.who.int/child-adolescent-health/New\\_Publications/NUTRITION/paper146.pdf](http://www.who.int/child-adolescent-health/New_Publications/NUTRITION/paper146.pdf)

Pan American Health Organization. *Guiding Principles for complementary feeding of the breastfed child*. Washington DC: Pan American Health Organization, World Health Organization, 2003.  
[http://www.who.int/child-adolescent-health/New\\_Publications/NUTRITION/guiding\\_principles.pdf](http://www.who.int/child-adolescent-health/New_Publications/NUTRITION/guiding_principles.pdf)

WHO. *Complementary feeding counselling: a training course*. WHO/NHD/CAH. Geneva: World Health Organization (in press)

WHO convened the Global Consultation on Complementary Feeding to update technical guidelines on complementary feeding and strengthen guidance on programmatic actions for their implementation. Background papers summarized state of the art knowledge on energy and nutrient requirements, factors affecting feeding behaviours, approaches for improving quality and utilization of complementary foods at household and central levels, and lessons learned from efforts to improve breastfeeding practices.

This report summarizes the conclusions and recommendations of the consultation. It includes revised guidelines for appropriate complementary feeding based on new calculations of energy and nutrient requirements, and accumulating evidence on the importance of responsive feeding. This information has been consolidated in Guiding Principles for Complementary Feeding of the Breastfed Child. The report also summarizes lessons learned from large-scale programmes to improve complementary feeding and proposes ten action steps that can be taken in health facilities to ensure that adequate support is provided for complementary feeding, along with support for sustained breastfeeding.

For further information please contact:

**Department of Child and Adolescent Health and Development (CAH)**

World Health Organization  
20 Avenue Appia  
1211 Geneva 27  
Switzerland

Tel +41-22 791 3281  
Fax +41-22 791 4853

website: <http://www.who.int/child-adolescent-health>


**Department of Nutrition for Health and Development (NHD)**

World Health Organization  
20 Avenue Appia  
1211 Geneva 27  
Switzerland

Tel: +41 22 791 3326  
Fax: +41 22 791 4156

website: <http://www.who.int/nut>

ISBN 92 4 154614 X



9 789241 546140

