

Index

- A/EEC *see* attaching and effacing *Escherichia coli*
- abattoir wastes 419–20
- abortions 234
- abrasions 33
- Acanthamoeba* 256, 259
- ascarid nematodes 295
- acceptable risks 24
 - see also* tolerable...
- acid-fast staining 285
- acquired immunodeficiency syndrome (AIDS)
 - cryptosporidiosis 142, 143
 - cyclosporiasis 283, 286
 - microsporidia 258, 326, 327
- acute gastrointestinal illness of unknown origin (AGI) 125
- adaptation capacities 9, 309, 310–11
- adenoviruses 343, 355
- aerobic digesters 414, 417
- aerosols 34, 38–9, 246
- AFLP *see* amplified fragment length polymorphism
- AFOs *see* animal feeding operations
- Africa 93–108
- age factors
 - animals 414
 - control envelope 79
 - Cryptosporidium* 183
 - dose-response studies 464
 - Giardia* 182–3
 - tolerable microbial risk 457–8
- AGI *see* acute gastrointestinal illness of unknown origin
- agriculture
 - animal systems 409–12
 - biosecurity 177

- campylobacteriosis risks 201
- industrial systems 21–4, 411
- mixed systems 411
- New Zealand 194, 196–7
- pastoral 194, 200
- public health impact 200
- Salmonella* outbreaks 231–2
- zoonosis distribution 52–3
- AIDS *see* acquired immunodeficiency syndrome
- air drying 413, 417
- Alaria marcianae* 300
- algal toxins 35
- alkaline treatment 414, 417
- alveolar hydatid disease 301–2
- ambient waters 442, 449, 450
- amitochondriate parasites 479
- amoebae 211, 256, 259–61
- Amoy Gardens, Hong Kong 38–9
- amplified fragment length polymorphism (AFLP) 352, 353
- anaerobic digestion 413, 415, 417
- Ancylostoma* 296
- Andean countries 307, 308
- Andes virus 105
- Angiostrongylus* 296–7
- animal densities 382
- animal faeces/wastes
 - Campylobacter* 176
 - Cyclospora cayetanensis* 287
 - cyclosporiasis 287
 - dead animals 419–21
 - disposal 22–3
 - Escherichia coli* O157 187–8
 - field application 416–17
 - human exposure 439–40, 443–4, 448–9
 - manure treatment 416
 - pathogen control 378–9, 409–22
 - pathogen sources 14
 - policies 418
 - Salmonella* 237
 - treatment processes 412–16, 417
 - versus human 450
- animal feeding operations (AFOs) 433–4
- animal husbandry
 - Campylobacter* 176–7
 - control envelope 69–71
 - emerging diseases 21–4
 - environmental impact 21–4
 - Escherichia coli* O157 178–9
 - global densities 22
 - global increase prediction 21
 - ‘livestock revolution’ 21
 - management 69–71
 - New Zealand 191–200
 - Salmonella* reservoirs 232–5
 - Scotland 167–8
 - stocking levels 199
 - zoonose distribution 52–3
 - see also* domestic animals
- animal products demand 6
- animal reservoirs
 - control strategies 377–8, 380–401
 - pathogen evaluation 9–10
 - risk factors 381–4
 - Salmonella* 232–5
 - Scotland 151, 167–8
 - viruses 246–50
- animal viral transmission 246
- anthropogenic factors 46–61
- anthroponoses 29
- antibiotic resistance analysis 352
- antibiotics
 - animal agriculture 412
 - Campylobacter jejuni* 71
 - control envelope 70, 71, 78
 - genetic exchange 477
 - resistance 71, 233, 238, 389–90, 477
 - Salmonella* 233, 238
 - virulence genes 78
 - zoonosis distribution 58
- antibodies 250, 251
- antigens 243

- antimicrobials 389–90
 Apicomplexa 265–75, 479
 ARCC *see* average rate of correct classification
 Argentina 105
Ascaris
 Asia/Africa/South America 94
 A. lumbricoides 122, 292, 294
 A. suum 294
 ascariis pneumonia 294
 Asia 93–108, 295
 asymptomatic infections 156, 158, 457
 at-risk groups
 control envelope 77
 future perspective 6
 microsporidia 326–8
 zoonosis distribution 55–6
 see also immunocompromised groups
 attaching and effacing *Escherichia coli* (A/EEC)
 214–15
 Australia 104, 231
 Australian bat lyssavirus 104
Austrobilharzia variglandis 122, 299
 average rate of correct classification (ARCC) 351
 average surface temperature 61
 avian influenza 104
 avian schistosomes 299
 avian viruses 248, 249

Bacillus anthracis 479
Bacillus coli *see* *Escherichia coli*
 backflow, distribution systems 434
 backwashing, filtration 427–8
 bacteria 210
 Asia/Africa/South America 101–3, 106
 control in animal wastes 411–12
 emergence/re-emergence 478–9
 expert consensus 12
 genetic exchange 477
 gorillas 101–2
 Mexico 115
 USA 123–4, 126, 132
 bacterial source tracking *see* faecal source
 identification; microbiological source
 tracking
 bacteriocins 392
 bacteriophages 343, 354, 392, 477
Bacteroides 355–6
 Bacteroidetes faecal markers 355–6, 359
Balantidium coli 121–2, 256
 bank filtration 433
 bats 103, 104–5
Baylisascaris procyonis 295
 beaches 339, 462
 behaviour, human 10, 53–4, 79–80
 Belgium 177
 benchmark organisms 8–9
 beta-Poisson models 464–5
 beverages 315–16
Bifidobacterium 357
 binning procedures 467
 biochips 374
 biofilms 55
 biosecurity 177, 388–9, 399–400
 biosensors 373–4
 birds 151, 167–8, 273
 black rat 310–11
Blastocystis 100, 121
Blastomyces hominis 116
 ‘boil water’ requirements 185, 187
 Bolivia 307, 308, 310
 bone meal 395, 398, 481
 bottled water 188
 bovine reservoirs 246–7
 bovine spongiform encephalopathy (BSE) 41–2,
 394–9, 400
 bovine reservoirs 247
 dead animal wastes 419–20, 421
 emergence/re-emergence 481
 Bradley’s classification 31–4

- broilers 22–3
- Brucella* 103
- BSE *see* bovine spongiform encephalopathy
- Bunyaviridae 105
- Burkholderia pseudomallei* 479
- Buxtonella* 256
- caffeine 350
- calciviruses 244, 248, 249
- calves and calving 171–2, 381, 390–1
- Campylobacter*
 - Asia/Africa/South America 102
 - farming transmission 222, 223
 - gastrointestinal illnesses 231
 - New Zealand 191–202
 - outbreak investigation 183–4
 - QRA application 461–3
 - risk models 158, 160
 - Scotland 169–70, 172, 176–7
 - seasonal patterns 171, 173
 - surveillance systems 163
 - survival rates 198
 - USA 123, 128–9, 132
 - C. coli* 102, 176–7, 371
 - C. jejuni*
 - Asia/Africa/South America 102
 - automated PCR detection 371
 - chlorine inactivation 429
 - control envelope 68, 71, 78
 - dose-response studies 464–5, 466
 - New Zealand 197
 - prevalence in animals 381
 - sarafloxacin 78
 - Scotland 176–7
- campylobacteriosis
 - case-study 461–3
 - causative agents 137
 - health consequences 138–9
 - management responses 201
 - New Zealand 192–3, 194, 197, 199
 - symptoms 137–8
 - treatment 138
- Canada 221, 222, 268
- Canary Islands 220
- candidate agents 246–50
- cantaloupes 231
- Capillaria hepatica* (syn. *Calodium hepaticum*) 297
- carbon-source utilization (CUP) 352
- Caribbean 113–18
- case definitions 159
- case-studies 151–207
- catchment basins 9–10
- cats 77, 267–9, 294
- cattle
 - BSE 394–9, 481
 - Campylobacter* 176
 - Cryptosporidium* 274
 - Escherichia coli* 51–2, 178–9, 210, 222, 223
 - industrialization 21–3
 - manure disposal 23
 - Mycobacterium avium* ssp. *paratuberculosis* 325
 - New Zealand 191–202
 - risk factors 381
 - Salmonella* reservoirs 233–4, 237
 - Salmonella* Typhimurium 172–3
 - Scotland 168, 171–3, 175
 - trade effects 51
 - viruses 246–7
 - VTEC reservoirs 217–18
- CDSC *see* Communicable Disease Surveillance Centre
- cell receptors 243, 245
- Central America 113–18
- cercarial dermatitis 299, 300
- cestodes *see* tapeworms
- chemical indicators 350, 447
- chickens *see* poultry
- children
 - congenital toxoplasmosis 145–8
 - diarrhoea in Latin America 114

- Dominican Republic 116
- El Salvador 117
- Giardia* infections 263
- Guatemala 118
- haemolytic uraemic syndrome 223
- Honduras 117
- Mexico 115
- mortality/morbidity 56–7
- Nicaragua 117
- tolerable microbial risk 457–8
- chips *see* microarray technology
- chloramine 432
- chlorine 74, 325, 384, 392–3, 429–31
- chlorine dioxide 432
- chronic wasting disease (CWD) 249, 252, 421, 481
- cisterns 236
- civets 38
- CLB *see* cyanobacteria-like bodies
- Clean Water Act* (US) 442–3
- climatic issues
 - climate change 58–61, 318
 - climate cycling 59
 - Cryptosporidium* outbreak 186
 - emerging zoonoses 477–8
 - fascioliasis transmission 317–18
 - greenhouse gases 418
- clinical parasitological studies 98–9
- cloned animals 444
- Clostridium perfringens* 369
- co-infections 29
- coagulation 427, 428
- coastal waters 236
- cold-tolerance 239
- coliforms 360, 431
- coliphages 343
- collaboration 164, 447
- colonization power 309, 312–13
- combining distributions 467
- Communicable Disease Surveillance Centre (CDSC) 163, 164
- community fingerprinting methods 357
- competitive exclusion 391–2
- composting 413, 415
- concentration distributions 467
- congenital toxoplasmosis 145–8
- conjugation 476
- Connecticut freshwater study 195–6
- consumption habits, human 53–4, 80
- contact diseases 32–3
- contamination
 - animal/human 450
 - beaches 339, 462
 - control envelope breach 75
 - Cryptosporidium* 274–5
 - epidemiological concerns 155
 - Giardia* 265
 - human faeces 194–5, 444
 - poultry 233
 - Stockholm framework 452–8
 - Toxoplasma* 269
- control envelope 66, 67–80
- control strategies
 - animal reservoirs 377–8, 380–401
 - animal wastes 378–9, 409–22
 - disease transmission 420
 - drinking-water 379, 426–35
 - future perspective 7
 - Listeria monocytogenes* 456
 - regulatory perspective 439–51
- Corsica 310–11
- Costa Rica 115–16
- coxsackieviruses 246
- Creutzfeldt-Jakob disease (CJD) *see* variant
 - Creutzfeldt-Jakob disease
- Crohn's disease 72, 189, 325
- cross-resistance, antibiotics 389–90
- cross-sector collaboration 16
- crossover between species 12
- cryptosporidiosis 256, 271

- Mexico 115
- New Zealand 192, 194, 195
- symptoms 142
- treatments 142–3
- Cryptosporidium* 269–75
 - age factors 183
 - Asia/Africa/South America 95–9
 - chlorine resistance 74
 - contamination reduction 274–5
 - control envelope breach 75
 - Cuba 116
 - disease models 159
 - HACCP-like processes 81
 - HIV 67–8
 - oocysts detection tests 96–8
 - outbreak investigation 183–4
 - regulation 441, 442
 - Scotland 169–70, 171–2, 180–1, 186–7
 - seasonal patterns 173
 - surface waters 97
 - surveillance 92, 163–4
 - treated waters 98
 - USA 121, 128–30, 131–2
 - UV light treatment 432–3
 - C. hominis* 180
 - C. parvum*
 - automated PCR detection 371
 - coagulation 427
 - control envelope 68
 - disease risk 339, 340
 - dose-response studies 464, 465
 - immunocompromised groups 384
 - immunoprophylaxis 390–1
 - prevalence in animals 381
 - resistance 390
 - surveillance 473–4
 - types 180
 - waste treatment systems 414
- Cuba 116
- culling 419, 420–1
- culture
 - bias 343–4, 354
 - Escherichia coli* O157:H7 218
 - culture-dependent faecal source identification 351–5
 - culture-independent faecal source identification 358
 - CUP *see* carbon-source utilization
 - CWD *see* chronic wasting disease
 - cyanobacteria-like bodies (CLB) 283
 - cyclophyllid cestodes 300–2
 - Cyclops* 297–8
 - Cyclospora cayetanensis* 116, 122, 211, 256, 283–8
 - cyclosporiasis 143, 283–8
 - cystic hydatid disease 301, 302
 - cysts 260, 262, 263–4, 266
- DAEC *see* diffuse adherent *Escherichia coli*
- dairy cattle
 - management responses 201
 - New Zealand 191–2, 194, 197, 198, 200, 201
 - public health impact 200
 - Scotland 168, 171
- DALYs *see* disability-adjusted life years
- dams 53
- DBPs *see* disinfection by-products
- dead-end hosts 28–9
- DEC *see* diarrhoeagenic *Escherichia coli*
- definition, zoonoses 5, 28
- deforestation 54
- demographic changes 55–7
- density of animals 382
- deposition/yield studies 199
- desiccation 413, 417
- design of distribution systems 434
- detection methods
 - Cryptosporidium* 96–8, 274
 - epidemiological surveillance 159
 - Giardia* cysts 96–8
 - microsporidia 258
 - outbreaks 161
 - rapid 367–74

- Salmonella* 237
- Toxoplasma* oocysts 268
- diarrhoea
 - age/income factors 458
 - cases per year 4
 - cryptosporidiosis 271
 - cyclosporiasis 284–5
 - giardiasis 263
 - Latin America 114
- diarrhoeagenic *Escherichia coli* (DEC) 213–24
- die-offs 420–1
- diffuse adherent *Escherichia coli* (DAEC) 214–15
- digenetic trematodes 292
- Diphyllobothrium* 303
- Diplogonoporus grandis* 303
- Diplostomum spathaceum* 300
- direct epifluorescent filter technique 370
- direct life cycles 292
- direct transmission 28–9
- direct zoonoses 28–9
- disability-adjusted life years (DALYs) 11, 454, 455–6, 457, 458
- discharge permits 449, 450
- disease emergence 19–25, 36–42, 339–40, 471–84
- disease models 157–8, 159
- disease specificity 230–1
- disease-based surveillance systems 159
- disinfectants 429–33
- disinfection by-products (DBPs) 428
- distribution systems 75, 434–5
- distribution of zoonoses
 - anthropogenic factors 46–61
 - calculation methods 467–8
 - Cryptosporidium* 271
 - Entamoeba histolytica* 260–1
 - environmental factors 46–61
 - Giardia* 262–3
 - human fascioliasis 307
 - microsporidia 257
 - Toxoplasma* 266
- DNA replication 476
- dogs 294
- domestic animals 217–18, 259, 267–9
 - see also* animal husbandry; pets
- domestic hygiene 32, 76, 77
- Dominican Republic 116–17
- dormancy 31
- dose-response studies
 - Campylobacter jejuni* 466
 - campylobacteriosis 462
 - regulation and control 448
 - uncertainties 464–5, 467
- Dracunculus medinensis* 32–3, 297–8
- drinking-water
 - Cyclospora cayetanensis* 287
 - diseases control 379, 426–35
 - emerging infection patterns 474
 - Entamoeba histolytica* 260
 - epidemiological concerns 155
 - Escherichia coli* O157:H7 222
 - faecal contamination 340–1, 342
 - indicators 342
 - regulation and control 275, 441–2, 449
 - risk factors 383
 - Stockholm framework 452–8
 - toxoplasmosis 268
 - USA outbreaks 128–9
- driving forces 5–6
- drought 59
- drug resistance 58, 107, 309
- drying animal waste 413, 417
- dynamic risk assessment 468
- EAggEC *see* enteroaggregative *Escherichia coli*
- Echinococcosis* 301–2
- education 77, 107–8
- effluents
 - Cryptosporidium* dispersal 272
 - health risks 200, 201
 - New Zealand 193, 194–5, 200, 201

- see also* animal faeces/wastes; human effluent/faeces
- EHEC *see* enterohaemorrhagic *Escherichia coli*
- EIEC *see* enteroinvasive *Escherichia coli*
- El Niño Southern Oscillation 59–60
- El Salvador 117
- emerging diseases 4, 19–25, 36–42, 339–40, 471–84
- empowerment 108
- Encephalitozoon* 57, 257–9, 326
- encephalomyocarditis virus 328, 329
- endemic diseases 161–2
- endogenous infections 213
- Enhanced Surface Water Treatment Rule (ESWTR) 275
- Entamoeba coli* 116
- Entamoeba histolytica* 260–1
- enteric organisms 228–39
- enteroaggregative *Escherichia coli* (EAaggEC) 214–15
- enterococci 342–3, 360
- Enterocytozoon bieneusi* 257–9, 326–8
- enterohaemorrhagic *Escherichia coli* (EHEC) 52–3, 210, 215, 339–40
- enteroinvasive *Escherichia coli* (EIEC) 214, 215
- enteropathogenic *Escherichia coli* (EPEC) 214, 215
- enterotoxigenic *Escherichia coli* (ETEC) 214, 215
- enterotoxins 356
- enteroviruses 243, 246–7, 343, 355
- enumeration methods 367–74
- environment introduction 9
- environmental changes 477–8
- environmental distribution factors 46–61
- environmental impact of animal agriculture 21–4
- environmental persistence 10, 30–1, 74
- Environmental Protection Agency (EPA) 441–4, 449–50, 457
- environmental resistance 10, 31, 74
- environmental transmission 30–1
- EPEC *see* enteropathogenic *Escherichia coli*
- epidemics, worldwide increase 46–7
- epidemiology
 - case-studies 151–207
 - concerns 155–9
 - Cryptosporidium* 273–4
 - cyclosporiasis 286–7
 - data 151–207
 - dynamic risk assessment 468
 - emerging disease patterns 20–1
 - human fascioliasis 308–9
 - limitations 154–5
 - New Zealand 194–6
 - outbreaks 151–207
 - protozoa 212
 - QRA relationship 460–1
 - regulatory approaches 443–4
 - risk management 81–2
 - Salmonella* 231–2
 - studies 154–65
 - surveillance 159–64, 170–1
 - VTEC 215–17
- Escherichia coli* 210
 - Asia/Africa/South America 102–3
 - Campylobacter* comparison 463
 - chlorine inactivation 429–30
 - concentrations 200
 - control envelope 74
 - deposition/yield studies 199
 - diarrhoeagenic 213–24
 - dose-response studies 464
 - faecal contamination indicator 336, 341, 342–3, 360
 - genetic diversification 57–8
 - health consequences 140
 - infectious dose 77
 - regulation 441–2
 - symptoms 139
 - toxin genes 356, 359
 - treatments 139
 - USA 123, 128–30, 132
- Escherichia coli* O6:H16 123, 128
- Escherichia coli* O121:H19 123, 130, 132
- Escherichia coli* O157
 - Asia/Africa/South America 102–3

- Scotland 167, 169–72, 177–9, 183–5, 187–8
- seasonal patterns 173
- Escherichia coli* O157:H7
 - agricultural practices 52–3
 - control envelope 68, 71–2, 73, 75
 - distribution system failures 434, 435
 - PCR detection 371
 - pH and survival 78
 - Swaziland outbreak 410
 - transmission 218–23
 - USA 123, 128, 130, 132
 - waste treatment systems 414, 415
- Escherichia coli* O157:[H7] 215–18
- estuarine areas 236
- ESWTR *see* Enhanced Surface Water Treatment Rule
- ETEC *see* enterotoxigenic *Escherichia coli*
- European surveillance systems 159
- excision 477
- exotic pets 85
- expert consensus 3–16
- exposure
 - calculation methods 467
 - control envelope 79
 - environmental transmission 31
 - epidemiological patterns 20–1
 - key concentrated sources 14
 - QRA 460–1
- eyes 295

- facultative lagoons/storage 413
- faecal coliforms 431
- faecal contamination
 - Bradley's classification 32
 - Escherichia coli* indicator 336, 341, 342–3
 - regulation 448–9
- faecal environmental loading 23–4
- faecal indicators 442–3, 446–7
- faecal source identification 349–61
 - see also* microbiological source tracking

- faecal wastes *see* animal faeces/wastes; human effluent/faeces
- farming *see* agriculture; animal husbandry
- Fasciola buski* 300
- Fasciola gigantica* 300, 305, 310, 312, 313
- Fasciola hepatica* 69, 299–300, 305, 310–13
- fascioliasis, human *see* human fascioliasis
- fasciolids
 - adaptation capacities 309, 310–11
 - colonization power 309
 - drug resistance 309
- fatty stools 263
- feedlots 389
- feral cats 269
- fields 416–17
- filtration
 - detection/enumeration methods 367–8
 - drinking-water 427–9
 - groundwater 433
 - parasites 479
- fingerprinting techniques 352–3, 357
- Finland 220
- fish kills 39, 40
- fish transmission 35
- flagellates 261–5
- flocculation 427
- flooding 59, 60, 324
- flukes 212
- flying fox 103
- Fonterra 201
- food
 - consumption habits 53–4, 80
 - epidemiological concerns 155
 - on-farm safety programmes 386–7
- food poisoning notification 170–1
- food processing effluent 237
- food sources
 - cyclosporiasis 286–8
 - hepatitis E 249
 - Listeria* 456

- Salmonella* 228–9, 231–2, 234–5
- Food Standards Agency 177
- forecasts 317–18
- Fossaria* 311, 312
- fowl plague 104
- fragrances 350
- France 327, 328
- Francisella tularensis* 124, 479
- freshwater plants 314–15
- freshwater quality criteria 442
- freshwater snail vectors 311–13
- freshwater studies
 - campylobacteriosis 462
 - Connecticut 195–6
 - Giardia* survival 264
 - New Zealand 196–8
- fungi 13
- Fung's Double Tube Method 369
- future trends 5–7, 471–84

- Galba truncatula* 311, 312–13
- gastrointestinal illnesses
 - Campylobacter* 195, 196, 197
 - child morbidity/mortality 56–7
 - Cryptosporidium* 273
 - Escherichia coli* 214
 - Giardia* 262
 - Salmonella* 231
 - surveillance systems 159–60
 - tolerable microbial risk 457–8
- genetic characterization 99
- genetic exchange 476–7
- genetic mutations 9, 57–8, 211, 475–6
- genetic variations 245
- genotypic methods 352–3
- genotyping 313, 326–7
- Giardia* 261–5
 - Asia/Africa/South America 95–9
 - chlorine resistance 74, 430–1
 - cyst detection tests 96–8
 - drinking-water regulations 275
 - filtration 428
 - regulation 441
- Scotland 169–70, 171–2, 181–3
- seasonal patterns 173
- surface waters 97
- treated waters 98
- USA 121, 128–30, 131–2
- UV treatment 432–3
- G. duodenalis* 95, 99, 381
- G. intestinalis* 457
- G. lamblia* 116, 144, 339, 340, 464
- giardiasis 144, 192, 193, 263
- Gigantobilharzia sturniae* 299
- global temperature rise 61
- globalization 51
- Gnathostoma spinigerum*, USA 123
- Gorilla gorilla* 101–2
- governments 440
- grazing systems 194, 196–200, 410–11
- 'great plate count anomaly' 343–4
- greenhouse gases 418
- groundwater 133, 236, 383, 433
- group benchmark pathogens 8–9
- Guangdong 37, 38
- Guatemala 118
- guidelines 452–8
- Guillian-Barré syndrome 138
- guinea worm 32–3, 297–8

- habitat creation 54
- HACCP *see* hazard analysis and critical control points
- haemolytic uraemic syndrome (HUS) 177, 178, 215–16, 223
- haemorrhagic fevers 105
- Haiti 286, 287
- hantavirus pulmonary syndrome (HPS) 105
- hantaviruses 105
- HAV *see* hepatitis A virus
- Hawaii 369
- hazard analysis and critical control points

- (HACCP) 80–1, 455
- hazard assessment, QRA 460–1
- health criteria, USA 448–50
- health education 107–8
- health policy, USA 444
- health targets 452–8
- heavy rainfall 60, 73, 382
- helminths 212, 291–303
 - cestodes 300–3
 - control in animal wastes 411–12, 415
 - expert consensus 13
 - life cycles 292
 - nematodes 293–8
 - transmission routes 292–3
 - trematodes 298–300
 - see also* parasites
- Hendra virus 103–4
- hepatitis A virus (HAV) 79, 124, 328, 329
- hepatitis E virus (HEV) 248, 249–50
 - control envelope 69
 - emergence/re-emergence 480
 - pigs 328–9
 - USA 124
- high-risk groups *see* at-risk groups; immunocompromised groups
- highly pathogenic avian influenza 104
- HIV *see* human immunodeficiency virus
- Honduras 117–18
- Hong Kong 38–9, 196, 339
- hookworms 292
- horizon-scanning perspectives 15
- host origin databases 351–3
- host specificity
 - control envelope 70
 - Salmonella* 230–1
 - viruses 243–4
- host susceptibility 10–11, 155, 156–7, 478
- host-agent-environment relationship 157–8
- host-specific polymerase chain reaction 359
- household factors, control envelope 76–7
- HPS *see* hantavirus pulmonary syndrome
- human activities impact 47, 48–9
- human behaviour 10, 53–4, 79–80
- human effluent/faeces
 - contamination 193, 194–5, 444
 - Cyclospora cayentanensis* 286
 - versus animal 201, 443, 450
- human fascioliasis 212, 305–18
 - adaptation/colonization power 309
 - environment colonization 317–18
 - epidemiology 308–9
 - geographical distribution 307
 - human infection sources 314–17
 - reservoir hosts 310–11
 - snail vectors 311–13
 - transmission 306–7
 - transmission foci 314
- human host changes 478
- human immunodeficiency virus (HIV) 257
 - Cryptosporidium* 67–8
 - cryptosporidiosis 142
 - Giardia* infections 263
 - toxoplasmosis 77
- human migration 10, 47–8, 54
- human population growth 6, 21, 49–50, 157
- human susceptibility 10–11
- human viral reservoirs 246
- human-to-human transmission *see* person-to-person transmission
- humans, *see also* pathogen–human factors
- Hurricane Mitch 117–18
- HUS *see* haemolytic uraemic syndrome
- hydatid disease 301–2
- hygiene 32, 76, 77
- ID *see* infectious dose
- IID *see* infectious intestinal disease
- immunity 11, 54, 76, 79
- immunocompromised groups 384
 - control envelope 79

- disease surveillance 84
- microsporidia 326, 327
- pathogen evaluation 11
- zoonose distribution 56
 - see also* at-risk groups
- immunoprophylaxis 390–1
- imported infections 182
- in vitro* infectivity 243–4
- income factors 6, 458
- index pathogens 454
- indicator methods
 - criteria requirements 446–7
 - EPA 450
 - limitations 444–5
 - viral indicators 343
 - water quality criteria 442–3
- indirect life cycles 292
- indirect transmission 21, 28, 29
- industrial farming systems 21–4, 411
- infants *see* children
- infection patterns, emerging 472–82
- infections with water-related insect vectors 34
- infectious agents environmental transmission 30–1
- infectious dose (ID) 31, 77
- infectious intestinal disease (IID) 183–4
- influenza viruses 248–9, 329
- infrastructure changes 54–5
- ingestion diseases 32–3
- inhalation 14, 34
- insect vectors 34
- Institute of Medicine of the National Academies (USA) 472–3
- interdisciplinary collaboration 16
- international issues
 - conflicts 50
 - control programmes 385–6
 - networks 85–6
 - surveillance 160–1, 164, 165
 - trade 85
 - travel 50–1
- invasive infections 213–14
- irrigation systems 53, 315
- Japan 219, 221, 295
- Johne's disease 72, 189, 324–5
- Juquitiba virus 105
- kidneys 243
- lagoons 413, 415
- lambling 172
- larva migrans 295
- latency 30
- laundry brighteners 350
- LEE *see* locus of enterocyte effacement
- Legionella* 34
- Legionnaires' disease 34, 54, 91
- length heterogeneity polymerase chain reaction (LH-PCR) 356, 357
- Leptospira* 124, 129–30, 132
 - L. canicola* 141
 - L. hebdomadis* 141
 - L. icterohaemorrhagiae* 141
 - L. interrogans*, USA 124
- leptospirosis 324
 - causative agents 141
 - climatic factors 478
 - Costa Rica 115
 - Honduras 117–18
 - Mexico 115
 - Scotland 169–70, 171
 - symptoms 141
 - transmission 33–4
 - treatments 142
 - vaccination 391
- LH-PCR *see* length heterogeneity polymerase chain reaction
- library-based faecal source identification 351–3
- life cycles
 - cestodes 292
 - Cryptosporidium* 270
 - Cyclospora cayatanensis* 284
 - Entamoeba histolytica* 260

- fasciolids 306–7
- Giardia* 262
- helminth parasites 292
- microsporidia 257
- nematodes 292
- Toxoplasma* 265–6
- trematodes 292
- zoonose criteria 35–6
- lime stabilization 414, 417
- Listeria* 371, 372, 456
- liver, raw 316
- liver fluke disease *see* human fascioliasis
- livestock *see* agriculture; animal husbandry
- lobose amoebae 259
- local communities empowerment 108
- local surveillance 160–1, 164
- locus of enterocyte effacement (LEE) 216
- Lyme disease 171
- lymnaeids 299, 311–13

- MAC *see* *Mycobacterium avium* complex
- ‘mad cow disease’ *see* bovine spongiform encephalopathy
- Malaysia 104–5
- malnutrition 11
- management
 - campylobacteriosis risks 201
 - distribution systems 435
 - future perspective 7
 - health risks 456
 - manure treatment 416
 - Stockholm framework 452–8
- management interventions, control envelope
 - human characteristics 79–80
 - livestock/animal management 70, 71
 - point of use/household 76, 77
 - source water protection 72–3
 - water supply 74–5
- MAP *see* *Mycobacterium avium* ssp. *paratuberculosis*

- MAR *see* multiple antibiotic resistance method
- marine studies 196, 264, 268–9
- marine water quality criteria 442
- mass culling 419, 420–1
- Maximum Contaminant Level Goals (MCLGs) 441–2
- MBM *see* meat and bone meal
- meat
 - consumption trends 21
 - demand 6
 - preparation 54
 - processing 85, 419–20
- meat and bone meal (MBM) 395, 398, 481
- membrane processes 428–9
- Menangle virus 103
- meningitis 297
- metacercariae 298
- Mexico 113–14, 115, 118
- microarray technology 342, 374
- microbial flora 235
- microbiological source tracking (MST) 447, 450, 451
 - see also* faecal source identification
- microfiltration membranes 429
- microsporidia 13, 95–9, 211, 256–9, 326–8
- microsporidiosis 257–8
- migrants, human 10, 47–8, 54
- military conflicts 50
- Miller Fisher syndrome 138
- Milwaukee, USA 440–1
- Ministry of Health (New Zealand) 202, 206–7
- mixed farming systems 411
- mobile genetic elements 477
- molecular beacon technology 372
- molecular techniques, protozoa 212
- molecular tracers 350
- monkeys 243
- Monte Carlo simulations 462, 467
- mortality
 - children 56–7
 - DALYs 455, 456

- foodborne listeriosis 456
 - helminth infections 291–2
- most probable number (MPN) methods 370
- mountain gorillas 101–2
- mouse studies 272
- movement
 - hosts 157, 164, 437, 478
 - pathogen reservoirs 47–51
 - pathogens 412
- MPN *see* most probable number methods
- MST *see* microbiological source tracking
- multibarrier protection 426–7
- multiple antibiotic resistance (MAR) method 352
- multiple array probe technologies 446
- multiple host susceptibility 292
- multiresistant strains 234, 238
- municipal water supplies 49
- mutation 9, 57–8, 211, 475–6
- mycobacterial aerosol transmission 34
- Mycobacterium avium* complex (MAC) 74, 479
- Mycobacterium avium* ssp. *paratuberculosis* (MAP) 68, 71, 72
- Myocastor coypus* 311

- Naegleria* 132, 256, 259
- nanofiltration membranes 428
- NASBA *see* nucleic acid sequence-based amplification technique
- national conflicts 50
- National Pollutant Discharge Elimination System (NPDES) 443, 450
- national surveillance 160–1, 164, 165
- Necator americanus* 296
- nematodes 212, 292, 293–8
- neonatal sepsis 146
- Nepal 286, 287
- Netherlands 23, 220
- New Zealand
 - beaches 339
 - campylobacteriosis 461–3
 - Ministry of Health pamphlet 206–7
 - risk models 158, 160
 - surveillance 152
 - zoonotic infections 191–202
- Nicaragua 117
- niche widening 312–13
- Nipah virus 104–5, 328, 329
- nitrate leaching 22
- non-O157 VTEC 216–17, 218
- non-VTEC DEC 224
- non-zoonotic agents 195
- noroviruses 184, 243, 480
- notifications 170–1
- NPDES *see* National Pollutant Discharge Elimination System
- nucleic acid segments 476
- nucleic acid sequence-based amplification technique (NASBA) 373
- nutria *see* *Myocastor coypus*
- nutrient pollution 418

- O157 VTEC 215–16
- occupational exposure 33, 170, 474
- ocular toxoplasmosis 145
- Office International des Epizooties (OIE) 385–7
- on-farm control 386–93
- Ontario 221, 222
- oocysts
 - Cryptosporidium* 181, 270, 272–3, 274
 - Cryptosporidium parvum* 473
 - Cyclospora cayetanensis* 284, 285
 - Toxoplasma* 265–8
- oral doses 233
- otters 268–9
- outbreaks
 - Campylobacter* 176
 - epidemiology 151–207
 - Scotland 183–7
 - surveillance systems 160, 162–3
- outcome factors 10–11
- oxidants 429–33
- ozonation 433

- pamphlets 202, 206–7
- pandemic influenza 328
- pandemic *Salmonella* 238
- paragonimiasis 35
- Paramyxoviridae 103, 104
- parasites
 - Asia/Africa/South America 94–100, 106
 - Cyclospora cayatanensis* 284, 287
 - emergence/re-emergence 479
 - see also helminths
- passive infected carriers 461–2
- pastoral farming 194, 196–200
 - see also animal husbandry
- pathogen-free animals 387
- pathogenicity islands 477
- pathogens
 - changes in 475–6
 - characteristics 57–8
 - control envelope 67, 68, 77–8
 - detection 341–2
 - emerging 4, 19–25, 36–42, 339–40, 471–84
 - main groups 12–13
 - reservoir mobility 47–51
- pathogen–human factors, control envelope
 - livestock/animal management 70–1
 - point of use/household 76
 - source water protection 71–2, 73
 - water supply 74
- patterns of infection 472–82
- PCR see polymerase chain reaction
- PEAS see possible estuary-associated syndrome
- pectin gel tubes 369
- persistence in environment 10, 30–1, 74
- person-to-person transmission
 - Escherichia coli* 210
 - Giardia* 183, 262
 - infectious intestinal disease 184
 - VTEC 223
- personal hygiene 32, 76, 77
- Peru 286, 287
- pets 76, 85, 235
- PFGE see pulsed-field gel electrophoresis
- Pfiesteria* 39–41
- pH 78
- phenotypes, new 475–6
- phenotypic methods 352
- picornaviruses 244, 245
- pigeons 235
- pigs
 - Campylobacter* 176
 - Fasciola hepatica* 311
 - manure disposal 22
 - production effects 52
 - Salmonella* reservoirs 234–5
 - Scotland 168
 - vesicular disease 246, 247
 - viruses 247–8, 328–9
 - waste lagoons 415
 - Yersinia enterocolitica* 210, 238–9
- pinniped caliciviruses 249
- plant ingestion 314–16
- plasmids 476–7
- Pocomoke River, Maryland 40
- point source exposure 20
- point of use 76–7
- policies
 - animal wastes 418
 - Listeria* control 456
 - risk profile usage 468
 - US public health 444
- polio 85–6, 430
- ‘pollution trading’ 449, 451
- polycystic echinococcosis 302
- polycystic hydatid disease 301
- polymerase chain reaction (PCR) 371–2
 - cyclosporiasis diagnosis 285
 - enteric virus quantification 355
 - methods 450
 - water quality surveillance 342, 344
- Pontiac fever 34

- population growth 6, 21, 49–50, 157
- possible estuary-associated syndrome (PEAS) 41
- potable water
 - acceptable risks 24
 - Asia/Africa/South America 106
 - Nicaragua 117
- poultry
 - biosecurity 388
 - Campylobacter* 71, 177
 - competitive exclusion 391–2
 - control programmes 386–7
 - industrialization 22
 - manure disposal 22–3
 - risk factors 381
 - Salmonella* 232–3
 - Scotland 168
 - undercooking 195, 196, 461
 - viruses 249
- precipitation
 - control envelope breach 73
 - global change 58–9
 - heavy rainfall 60, 73, 382
- pregnancy 77, 145–6, 266
- presence/absence data 468
- pressure, water systems 434, 435
- prevalence
 - Cryptosporidium* 271
 - Entamoeba histolytica* 260–1
 - geographical factors 157
 - Giardia* 262–3
 - microsporidia 257
 - socioeconomic factors 157
 - Toxoplasma* 266, 267
- preventative quality assurance framework 83
- primers 356, 358
- prions 12, 394–9, 421, 481
- private water supplies 168–9, 185, 474
- probability of illness 465
- probiotics 392
- processing methods 85
- protein coating 242–3, 244, 245
- protozoa 211–12, 255–76
 - expert consensus 13
 - transmission routes 95
 - USA 121–3, 126, 131–2
- pseudophyllid cestodes 300, 303
- Pteropus* 103, 104
- public health
 - agriculture impact 200
 - driving forces 5–6
 - pathogen evaluation 11–12
 - Stockholm framework 452–8
- pulsed-field gel electrophoresis (PFGE) 352, 353
- QRA *see* quantitative risk assessment
- quality assurance 83
- quality criteria
 - ambient waters 449, 450
 - Clean Water Act* US EPA 442–3
 - water distribution systems 435
- quantitative microbial risk assessment (QMRA) 81–2
- quantitative risk assessment (QRA) 460–8
- quarantine procedures 70, 71
- racehorses 103
- racoons 295
- Radix* 312
- rainfall *see* precipitation
- rapid detection/enumeration methods 367–74
- rats 250, 310–11
- raw liver 316
- re-emergence 4, 478–81
- reassortant virus strains 29
- recombination of genetic elements 477
- recontamination 55
- recreational waters 33
 - campylobacteriosis 461–2
 - Cryptosporidium* 181
 - emerging patterns 474
 - epidemiological concerns 155
 - Escherichia coli* O157:H7 218, 222
 - faecal contamination 340–1

- faecal indicators 342
- New Zealand 193–6, 197–9
- quality criteria 442, 443
- regulatory approaches 441
- risk factors 383
- swimming-associated illness 339, 341
- USA 127, 129–30, 133
- regional effects 59
- regulations
 - Cryptosporidium* contamination 275
 - Toxoplasma* contamination 269
- regulatory perspective 437–51
- Reiter's syndrome 138
- remote sensing 318
- repetitive extragenic palindromic polymerase chain reaction (REP-PCR) 352, 353
- replication 31, 244, 476
- reservoirs
 - animal 377–8, 380–401
 - human fascioliasis 310–11
 - Salmonella* 232–5, 237
 - Scotland 167
 - viruses 246–50
 - VTEC 217–18
- resistance
 - antibiotics 58, 389–90
 - chlorination 384
 - drugs 58, 107, 309, 389–90
 - environmental 10, 31, 74
 - environmental stressors 9
- respiratory illnesses 196
- reverse osmosis membranes 428
- ribonucleic acid (RNA) 57, 244, 480
- ribotyping 352, 353
- risk assessment 437–69
 - emerging zoonotic diseases 24–5
 - modelling 153, 156, 161
 - quantitative 460–8
 - Stockholm framework 452–8
 - uses 437–8
 - water usage 441
- risk communication 77
- risk factors
 - animal reservoirs 381–4
 - diseases occurrence/spread 47
- risk management 66, 80–6
 - disease surveillance 84–5
 - epidemiological studies 81–2
 - HACCP system 80–1
 - international networks 85–6
 - QMRA 81–2
 - quality surveillance 338–44
 - targeted studies 84–5
 - water safety plans 82–4
- risk profiles 461, 462–3, 467–8
- rivers *see* streams and rivers
- RNA *see* ribonucleic acid
- rodent reservoirs 250
- rotaviruses 247, 480
- ruminants
 - biosecurity 388–9
 - Cryptosporidium* infections 270
 - risk factors 381
 - see also* cattle; sheep
- runoff 22, 73
- rural health regions 194–5, 197, 200
- Safe Drinking Water Act* (US) 441–2
- SALINPORK programme 387
- Salmonella* 210, 228–39
 - Asia/Africa/South America 101–2
 - competitive exclusion 391, 392
 - definition 229
 - dose-response studies 464, 465
 - emerging problems 238
 - Mexico 115
 - outbreak investigation 183, 184
 - pathogen-free animals 387
 - PCR detection 371, 372
 - Scotland 169–70, 172–5

- seasonal patterns 171, 173
- USA 123, 128–9, 132
- S. Abortusovis* 234
- S. bongori* 230
- S. Choleraesuis* 234–5
- S. enterica* 229, 230, 238, 381
- S. Enteritidis* 172, 174, 232–3, 238
- S. Kottbus* 231
- S. Poona* 231
- S. Typhimurium* 172–4, 232, 233–4, 238
- salmonellosis 140–1, 192
- sanitary surveys 435
- sanitation, control envelope 76, 77
- sapoviruses 243
- sarafloxacin-susceptible/resistant strains 78
- SARS *see* severe acute respiratory syndrome
- SCCWRP *see* Southern California Coastal Water Research Project
- Schistosoma* 33, 298–9
- Schistosomatidae 129–30
- schistosome cercarial dermatitis, USA 122
- Scotland 151, 167–89, 219–21
- scrapie 395, 397, 398–9, 481
- sea level rise 59
- sea lions 249
- sea mammals 264, 268–9
- seafood *see* shellfish
- seals 264
- seasonality 59–60
 - Campylobacter* 197, 198–9
 - Cryptosporidium* 164
 - human fascioliasis 317
 - Scotland 171–2, 173
- seawater
 - Cryptosporidium* 272
 - Entamoeba histolytica* 260, 261
 - Giardia* 264
 - microsporidia 259
- secondary transmission 155, 156
- sedimentation, drinking-water 427
- sediments 60, 199
- self-contained film system 369
- sensitivity, surveillance systems 159, 161, 162
- Seoul virus 105
- severe acute respiratory syndrome (SARS) 37–9, 105–6, 246
 - control envelope 77
 - emergence/re-emergence 480
 - host change capacity 57
- sewage *see* animal faeces/wastes; effluents; human effluent/faeces
- shedding 246, 262
- sheep
 - Campylobacter* 176
 - Cryptosporidium* 186–7, 274
 - MAP-infected 72
 - New Zealand 191
 - Salmonella* 234
 - Scotland 168, 172
 - scrapie 395, 398–9, 481
- shellfish
 - Campylobacter* 198, 462
 - growing and harvesting 443
 - transmission 35
 - transportation 51, 54
 - water quality criteria 443
- shellfish waters
 - animal/human faeces 340–1
 - faecal indicators 342
- Shiga toxin-producing *Escherichia coli* (STEC) 192, 194, 210, 214, 219–21
- Shigella* 101–2, 215
- skin injuries 33
- snails
 - cercarial dermatitis 299
 - Fasciola hepatica* 299
 - Galba truncatula* 311, 312–13
 - human fascioliasis 311–13
 - lymnaeids 299, 311–13
 - Schistosoma* 298
- socioeconomic factors 157
- sodium chlorate 393

- soils, animal wastes application 416
- sorbitol 216
- source tracking 349–61, 447, 450, 451
- source water protection 71–3
- South Africa 219
- South America 93–108
- Southern California Coastal Water Research Project (SCCWRP) 358–60
- specific pathogen-free animals 387
- spill over exposure 21
- spiral plating method 368
- Spirochetes emergence/re-emergence 479
- Spirometra* 122–3, 303
- sporadic diseases 161–2
- spores 257, 258–9
- sporulation 265–6, 267–8, 270
- sprouts 231–2
- square bacteriological filter 368–9
- stability, viruses 246
- standards, US 448–50
- statutory reporting 170–1
- steatorrhoea (fatty stools) 263
- STEC *see* Shiga toxin-producing *Escherichia coli*
- Stockholm framework 452–8
- stocking levels 199
- stomach acidity 210
- stool samples 163, 175
- storage, animal wastes treatment 413
- storm events 73, 74
- strains
 - differentiation 15
 - dose-response uncertainties 465
 - infectious dose variation 77
- streams and rivers
 - Campylobacter* 198–9
 - Escherichia coli* 200
 - management 201
 - sampling 197
 - sediments 199
- streptococci 360
- Strongyloides* 100, 296
- surface temperature 61
- surface waters
 - Entamoeba histolytica* 260
 - grazing systems 410
 - microsporidia 258–9
 - Salmonella* transmission 236
- surveillance
 - diseases outbreaks 162–3
 - Dominican Republic 116–17
 - emerging pathogens 473–4
 - epidemiological 159–64
 - importance 91–2, 152–3
 - limitations 153, 154–5
 - primary purposes 160–1
 - QRA relationship 460–1
 - risk management 84–5, 338–44
 - Scotland 170–88
 - studies 154–65
 - US 120–1
 - veterinary 163–4
 - water quality 338–44
- survival rates 31
 - Campylobacter* 198
 - Cryptosporidium* 272
 - Giardia* cysts 263–4
 - microsporidia 258–9
 - Toxoplasma* 267–8
- susceptibility
 - changes 155, 156–7, 158, 478
 - pathogen evaluation 10–11
- Swaziland 410
- Sweden 163
- swimmer's itch *see* schistosome cercarial dermatitis
- swimming pools *see* recreational waters
- swine *see* pigs
- symptom-based surveillance systems 159–60
- system failures 82
- systematic errors 156

- T-RFLP *see* terminal restriction fragment length polymorphisms
- Taenia* 122, 301
- tapeworms 122, 212, 292, 300–3
- targeted studies 84–5
- technological interventions, control envelope
 - human characteristics 79–80
 - livestock/animal management 70, 71
 - point of use/household 76, 77
 - source water protection 72–3
 - water supply 74–5
- temperature rise 61
- terminal restriction fragment length polymorphisms (T-RFLP) 356, 357
- thermophilic *Campylobacter* 198
- thermotolerance 229
- tiered monitoring 447, 449
- Tioman virus 103
- TMDLs *see* total maximum daily loads
- tolerable disease burden 83–4
- tolerable risks 454, 456–8
- total maximum daily loads (TMDLs) 449, 450, 451
- total viable cell count method 368–9
- tourism 50–1
- toxin genes 356, 359
- Toxocara* 294–5
- Toxoplasma* 99–100, 265–9
- Toxoplasma gondii* 68, 77, 122, 144–5
- toxoplasmosis 266, 268
 - congenital 145–8
 - health consequences 148
 - ocular 145
 - Scotland 171
 - symptoms 144–5
 - treatments 146–8
- trade issues 51, 85
- traditional livestock systems 21
- training 107–8
- transboundary movements 437
- transformation process 476
- transmissible spongiform encephalopathy (TSE) 394–9, 421, 481
- transmission
 - Bradley's classification 31–4
 - control 420
 - criteria 36
 - Cryptosporidium* 270
 - cyclosporiasis 286–7
 - emerging diseases 21
 - emerging patterns 474
 - Entamoeba histolytica* 260
 - environmental 30–1
 - fish 35
 - foci 314
 - Giardia* 262
 - helminths 292–3
 - human fascioliasis 306–7, 314
 - microsporidia 257
 - modes 5, 155–6, 158
 - multiple modes 5
 - principle vehicles 14
 - protozoan parasites 95
 - shellfish 35
 - Toxoplasma* 266–7
 - TSE 481
 - water aerosols 34
- transport
 - animal wastes 412, 416, 420
 - shellfish 51, 54
- travel issues 50–1, 85
- treatment
 - amoebiasis 261
 - animal wastes 412–16, 417
 - cryptosporidiosis 271
 - cyclosporiasis 285
 - drinking-water 426–34
 - Giardia* 263
 - microsporidiosis 257–8
- trematodes 292, 298–30
- Trichobilharzia* 299
- Trichuris* 122, 123, 292, 295

- triclabendazole 309
- tropical forest invasion 54
- tropical organisms 93–108
- tropism 243
- TSE *see* transmissible spongiform encephalopathy
- tularaemia 124
- typhoid 192
- typing schemes 176, 177, 180

- UDG *see* uracil-D-glycosylase
- Uganda 101–2
- ultrafiltration membranes 429
- ultraviolet (UV) light treatment 432–3
- United Kingdom
 - Escherichia coli* O157 isolates 178, 179
 - Salmonella* 233
 - Scotland 151, 167–89, 219–21
 - surveillance systems 163
 - VTEC 218, 219–21, 222
- United States of America 120–34
 - beaches 339
 - control programmes 152
 - cryptosporidiosis 440–1
 - Cryptosporidium* 273
 - Cyclospora cayetanensis* 286, 287
 - distribution systems 434
 - drinking-water regulations 275
 - Giardia* 262–3
 - health criteria 448–50
 - hepatitis E virus 329
 - Institute of Medicine of the National Academies 472–3
 - Listeria* control 456
 - microsporidia 327–8
 - principal pathogens 121–5
 - regulatory approaches 441–5
 - reported outbreaks 125–30
 - Salmonella* 231
 - standards 448–50
 - surveillance systems 163
 - tolerable microbial risks 457
 - Toxoplasma* 267, 268
 - VTEC outbreaks 219–21, 222
- uracil-D-glycosylase (UDG) 372
- urban agriculture 6
- urban wildlife 6
- urbanization 6, 49–50
- urine 23
- uropathogenic *Escherichia coli* 213, 214
- UV light *see* ultraviolet light

- vaccination
 - control envelope 76
 - immunoprophylaxis 390–1
 - wildlife 393–4
 - worm infection effects 94
- variant Creutzfeldt-Jakob disease (vCJD) 42, 247
 - emergence/re-emergence 481
 - potential waterborne pathogens 396, 397, 398, 400
- vegetables 286, 287–8
- verocytotoxin-producing *Escherichia coli* (VTEC) 192, 194, 210, 213–24
- vertical transmission 232–3
- vesicular diseases 246, 247, 249
- veterinary surveillance 163–4
- viable but non-culturable (VBNC) state 31, 198, 237
- Vibrionaceae 479
- Vietnam 286, 287
- virulence
 - control envelope 77–8
 - Escherichia coli* 215
 - Salmonella* 230–1
- viruses 211, 242–52
 - Asia/Africa/South America 103–6
 - control in animal wastes 411–12
 - drinking-water regulations 275
 - emergence/re-emergence 480

- expert consensus 12
- faecal source identification 354, 355
- faecal virus indication 343
- host specificity 243–4
- Mexico 115
- pigs 328–9
- replication 244
- USA 124–5, 126
- vital dyes 370
- VTEC *see* verocytotoxin-producing *Escherichia coli*

- Walkerton, Ontario 222
- washing 316
- Washington County Fair 222
- wastes *see* animal faeces/wastes; effluents; human effluent/faeces
- water distribution, control envelope 73–5
- water quality
 - assurance framework 81
 - criteria 435, 442–3, 449, 450
 - management 82–3
 - targets 83
- water safety plans (WSP) 82–4
- water shortages 93, 106
- water storage, control envelope 73–5
- water systems infrastructure changes 54–5
- water treatment
 - control envelope 73–5
 - emerging patterns of infection 474
 - parasites 479
 - viruses 480
- water-based infection classification 32–3
- water-borne infection classification 32
- water-related disease
 - classification 36, 37
 - criteria 30–5
- water-washed infections 32
- waterborne outbreaks definition 120
- watercress 314–15
- weather patterns 59–60

- whipworms *see* *Trichuris*
- WHO *see* World Health Organization
- wildlife
 - microsporidia 259
 - reservoirs 389, 393–4, 400
 - SARS virus 38
 - Toxoplasma* prevalence 267
 - viruses 249–50
 - VTEC reservoirs 217–18
- Wisconsin, USA 440–1
- World Health Organization (WHO) 452–8
- worm infections 32–3, 94
- wounds 33
- WSP *see* water safety plans

- years of life lost to premature death (YLL) 455
- years lived with a disability (YLD) 455
- Yersinia* 123–4, 128–9
- Yersinia enterocolitica* 123–4, 210, 238–9, 339
- yield studies *see* deposition/yield studies
- young animals 381

- zoonoses
 - definition 5, 28
 - determining criteria 28–9